

# LAMPIRAN

## Lampiran 1. Analisa Data

### Usia

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid >= 40 th	38	54,3	54,3	54,3
< 40 th	32	45,7	45,7	100,0
Total	70	100,0	100,0	

### Hipertensi

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Ya	29	41,4	41,4	41,4
Tidak	41	58,6	58,6	100,0
Total	70	100,0	100,0	

### Penyakit Kardiovaskular

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Ya	30	42,9	42,9	42,9
Tidak	40	57,1	57,1	100,0
Total	70	100,0	100,0	

### Trauma

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Ya	8	11,4	11,4	11,4
Tidak	62	88,6	88,6	100,0
Total	70	100,0	100,0	

### Idiopatik

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Ya	7	10,0	10,0	10,0
Tidak	63	90,0	90,0	100,0
Total	70	100,0	100,0	

### Epistaksis

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Posterior	35	50,0	50,0	50,0
Non Posterior	35	50,0	50,0	100,0
Total	70	100,0	100,0	

## Crosstabs Usia \* Epistaksis

Crosstab

			Epistaksis		Total
			Posterior	Non Posterior	
Usia	>= 40 th	Count	26	12	38
		% within Usia	68,4%	31,6%	100,0%
		% within Epistaksis	74,3%	34,3%	54,3%
		% of Total	37,1%	17,1%	54,3%
	< 40 th	Count	9	23	32
		% within Usia	28,1%	71,9%	100,0%
		% within Epistaksis	25,7%	65,7%	45,7%
		% of Total	12,9%	32,9%	45,7%
Total		Count	35	35	70
		% within Usia	50,0%	50,0%	100,0%
		% within Epistaksis	100,0%	100,0%	100,0%
		% of Total	50,0%	50,0%	100,0%

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	11,283 <sup>b</sup>	1	,001		
Continuity Correction <sup>a</sup>	9,729	1	,002		
Likelihood Ratio	11,619	1	,001		
Fisher's Exact Test				,002	,001
Linear-by-Linear Association	11,122	1	,001		
N of Valid Cases	70				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 16,00.

## Hipertensi \* Epistaksis

Crosstab

			Epistaksis		Total
			Posterior	Non Posterior	
Hipertensi	Ya	Count	20	9	29
		% within Hipertensi	69,0%	31,0%	100,0%
		% within Epistaksis	57,1%	25,7%	41,4%
		% of Total	28,6%	12,9%	41,4%
	Tidak	Count	15	26	41
		% within Hipertensi	36,6%	63,4%	100,0%
		% within Epistaksis	42,9%	74,3%	58,6%
		% of Total	21,4%	37,1%	58,6%
Total	Count	35	35	70	
	% within Hipertensi	50,0%	50,0%	100,0%	
	% within Epistaksis	100,0%	100,0%	100,0%	
	% of Total	50,0%	50,0%	100,0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	7,124 <sup>b</sup>	1	,008		
Continuity Correction <sup>a</sup>	5,887	1	,015		
Likelihood Ratio	7,266	1	,007		
Fisher's Exact Test				,015	,007
Linear-by-Linear Association	7,022	1	,008		
N of Valid Cases	70				

a. Computed only for a 2x2 table

b. 0 cells (,0%) have expected count less than 5. The minimum expected count is 14,50.