

## INTISARI

**Latar belakang:** Seseorang yang hidup di dataran tinggi dan di dataran rendah akan mengadakan adaptasi respirasi yang berbeda. Tekanan atmosfir, tekanan parsial oksigen, dan saturasi oksigen lebih rendah pada dataran tinggi. Hal tersebut dapat mempengaruhi tes fungsi paru pada penduduk yang telah lama tinggal di dataran tinggi karena tubuh beradaptasi untuk menyesuaikan keadaan sekitar. **Tujuan:** Mengetahui adanya perbedaan hasil FVC, FEV<sub>1</sub>, dan rasio FEV<sub>1</sub>/FVC pada penduduk yang tinggal pada ketinggian berbeda yaitu di dataran tinggi Kaliurang (875 m dpl) dan dataran rendah pantai Parangtritis (100 m dpl). **Bahan dan Cara:** Rancangan penelitian ini menggunakan jenis penelitian observasi analitik dengan pendekatan *cross-sectional* (potong lintang). Total 60 subyek siswa SMAN 1 Kretek, SMK Muhammadiyah Kretek Bantul, dan SMK Hamong Putera telah diteliti untuk tes fungsi paru dinamik. FVC dan FEV<sub>1</sub> diukur dengan spirometer manual. Analisis statistik yang dipakai adalah *Mann-Whitney* dan *Independent sample t – test*. **Hasil:** FVC, FEV<sub>1</sub>, rasio FEV<sub>1</sub>/FVC dianalisis dan dibandingkan antara kedua kelompok. Pada penelitian ini, FVC dan FEV<sub>1</sub> kelompok dataran tinggi Kaliurang dengan kelompok dataran rendah Parangtritis tidak terdapat perbedaan yang signifikan dengan nilai koefisien berturut-turut 0,219 dan 0,095, sedangkan rasio FEV<sub>1</sub>/FVC kelompok subyek dataran tinggi dan kelompok subyek dataran rendah berbeda signifikan ( $p=0,001$ ). **Kesimpulan:** perbedaan tempat tinggal akan menyebabkan perbedaan rasio FEV<sub>1</sub>/FVC akibat dari adaptasi lingkungan dan rasio FEV<sub>1</sub>/FVC penduduk dataran tinggi lebih tinggi daripada penduduk dataran rendah dengan hasil rerata rasio FEV<sub>1</sub>/FVC pada dataran tinggi Kaliurang 91% dan dataran rendah Parangtritis 86%.

**Kata kunci:** dataran tinggi, dataran rendah, tes fungsi paru, FVC, FEV<sub>1</sub>

## ABSTRACT

**Background:** People who live in high altitude and low altitude will hold different respiratory adaptation. The atmospheric pressure, oxygen partial pressure, and oxygen saturation is lower in highland. It can influence lung function test on people who have been living in high altitude for a long time because the body will adapt to adjust the surrounding circumstances. **Aim:** Determining the difference of FVC, FEV<sub>1</sub>, and FEV<sub>1</sub>/FVC ratio results on people who live in difference heights; which is in Kaliurang high altitude (875m above sea level) and Parangtritis low altitude (100m above sea level). **Materials and Methods:** This research uses analitic observational study with cross-sectional. The total of 60 subjects from students in SMAN 1 Kretek, SMK Muhammadiyah Kretek Bantul, and SMK Hamong Putera, Pakem, Sleman were assessed for dynamic lungs function test. FVC and FEV<sub>1</sub> is measured by manual spirometer. Statistical analyzing in this research uses Mann-Whitney and Independent sampel t-test. **Result:** FVC, FEV<sub>1</sub>, rasio FEV<sub>1</sub>/FVC were analyzed and compared between two group. In this research FVC and FEV<sub>1</sub> on the Kaliurang highland group is no significantly different with Parangtritis lowland group, the p value are 0,219 and 0,095, while the ratio of FEV<sub>1</sub>/FVC group in highland and lowland is significantly different ( $p=0,001$ ). **Conclusion:** The difference of geographical location will affect the different FEV<sub>1</sub>/FVC ratio that is caused by environment adaptation and the FEV<sub>1</sub>/FVC ratio on the people who live in highland is higher than the people who live in lowland. The mean of FEV<sub>1</sub>/FVC ratio on high altitude person is 91% while on low altitude person is 86%

**Keywords:** highland, lowland, lung funtion test, FVC, FEV<sub>1</sub>