

DAFTAR PUSTAKA

- Ali, I. (2006). *Mengatasi Gangguan Pada Telinga Dengan Tanaman Obat*. Jakarta: Agro Media Pustaka.
- Al-Qur'an. Surat Al-An'am ayat 46.
- Bielecki, I., Horbulewicz, A., Wolan, T. (2011). Risk factor associated with hearing loss in infants: An analysis of 5282 referred neonates. *Int J Ped Otorhinolaryngol*, 75, 925-30.
- Cambridge Communication Limited (1999). *Anatomi Fisiologi : Sistem Lokomotor Dan Penginderaan, Ed.2*. (Asih, Y., penerjemah). Jakarta: EGC. (Buku asli diterbitkan 1977).
- Chandra, B. (1996). *Pengantar Prinsip dan Metode Epidemiologi*. Jakarta: EGC.
- Cloherty, P.J., Eichenwald, E.C., Hansen, A.R., Stark, A.R. (2012). *Manual Of Neonatal Care, Ed.7*. Philadelphia: Lippincott Williams & Wilkins.
- Daniel, S.J., Waissbluth, S., Akinpelu, O.V. (2013). *Auditory risk of hyperbilirubinemia in term newborns: A systematic review*. South Carolina : The Society for Ear, Nose and Throat Advances in Children
- Direktorat Jenderal Bina Pelayanan Medik Kementerian Kesehatan RI. (2004). *Health Technology Assessment*(Tatalaksana Ikterus Neonatorum). Jakarta.
- Direktorat Jenderal Bina Pelayanan Medik Kementerian Kesehatan RI. (2010). *Health Technology Assessment*(Buku Panduan Tatalaksana Bayi Baru Lahir Di Rumah Sakit). Jakarta.
- Gorga, M.P., et al. (2000). Identification of neonatal hearing impairment: distortion product otoacoustic emissions during the perinatal period. *Ear Hearing*, 21.400-24.
- Hearing, J.C.o.I. (2007). Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs. *Pediatrics*, 120(4), 898-921.
- Herwanto, R.Y. (2012). *Gambaran gangguan pendengaran pada bayi dengan hiperbilirubinemia di RSUP H Adam Malik Medan [abstrak]*. Karya Tulis Ilmiah strata dua, Universitas Sumatera Utara , Medan.
- Isselbacher, K.J., Braunwald, E., Wilson, J.D., Martin, J.B., Fauci, A.S., Kasper, D.L. (1999). *Harrison Prinsip-Prinsip Ilmu Penyakit Dalam, Ed.13*,

Vol.1. (Asdie, A., H., penerjemah). Jakarta: EGC. (Buku asli diterbitkan 1994).

- Johnson, J.L., White, K.R., Widen, J.E., Gravel, J.S., James, M., Kennalley, T., Holstrum, J. (2005). A Multicenter Evaluation of How Many Infants With Permanent Hearing Loss Pass a Two-Stage Otoacoustic Emissions/Automated Auditory Brainstem Response Newborn Hearing Screening Protocol. *Pediatrics*, 116(3), 663-672.
- Lilyana. (2008). *Faktor-faktor risiko hipertensi pada jamaah pengajian majelis dzikir sby nurussalam tahun 2008*. Skripsi strata satu, -Universitas Indonesia, Depok.
- Mansjoer, A., Suprohaita, Wardhani, W.I., Setiowulan, W. (2009). *Kapita Selekta Kedokteran*. Jakarta: Media Aesculapius Fakultas Kedokteran UI.
- Martinez-cruz, C.F., Poblano, A., Fernandez-Carrocer, L.A. (2008). Risk factors associated with sensorineural hearing loss in infants at the neonatal intensive care unit: 15-year experience at the national institute of perinatology (Mexico city). *Arch Med Res*, 39, 686-94.
- Murti, B. (2003). *Prinsip dan Metode Riset Epidemiologi*. Yogyakarta: Gajah MadaUniversity Press.
- Northern, J.L., Downs, M.P. (2002). *Hearing In Children, Ed.5*. Pennsylvania: Lippincott Williams & Wilkins.
- Sarosa, G.I., Putranti, A.H., Susanto, J.C. (2010). Risiko Gangguan Pendengaran pada Neonatus Hiperbilirubinemia. *Sari Pediatri*, 12(4), 222-7.
- Sarosa, G.I., Putranti, A.H., Susanto, J.C. (2010). Risiko Gangguan Pendengaran pada Neonatus Hiperbilirubinemia [abstrak].
- Sarosa, G.I., Putranti, A.H., Setyarini, T.K. (2011). Pengaruh Asfiksia Neonatal Terhadap Gangguan Pendengaran. *Sari Pediatri*, 13(1), 5-13.
- Saricci, S.U., Serdar, M.A., Korkmaz, A., *et al.* (2004). Incidence, course, and prediction of hyperbilirubinemia in near-term and term newborns. *Pediatrics*. pp.775-80.
- Wong, D.L. (2003). *Pedoman Klinis Keperawatan Pediatrik, Ed.4*. (Ester, M., penerjemah). Jakarta: EGC. (Buku asli diterbitkan 2000).
- Wong, D.L. (2008). *Buku Ajar Keperawatan Pediatrik, Ed.6, Vol.1.* (Sutarna, A., penerjemah). Jakarta: EGC. (Buku asli diterbitkan 2001).