

DAFTAR PUSTAKA

- Aini, Dwi Nur. (2012). Pengaruh Latihan Fisik Terhadap Fatigue Pada Pasien Intrahemodialisis Di RSUD Tugurejo Semarang. Tesis.
- Applicability, Validity, and Reliability of the Piper Fatigue Scale in Postpolio Patients.* Available from: https://www.researchgate.net/publication/232163617_Applicability_Vailidity_and_Reliability_of_the_Piper_Fatigue_Scale_in_Postpolio_Patients [accessed Jan 22, 2016]
- Asri.P. Marthon, Marjono, Purwanto. (2006). Hubungan Dukungan sosial dengan tingkat depresi pasien yang menjalani terapi hemodialisis. Diakses 15 November 2016. <http://www.i-lib.ugm.ac.id>
- Astroth, K. S., Russell, C. L., & Welch, J. L. (2013). Non-pharmaceutical fatigue interventions in adults receiving hemodialysis: A systematic review. *Nephrology Nursing Journal*, 40(5), 407-27; quiz 428. Retrieved from <http://search.proquest.com/docview/1444878904?accountid=38628>
- Ayu, G.I., (2010). Hubungan Antara quick of blood (Qb) dengan adekuasi hemodialisis pada pasien yang menjalani terapi HD di Ruang HD BRSU Daerah Tabanan Bali, Tesis.
- Azwar, S. (2005). *Sikap Manusia dan Pengukurannya*. Jakarta: Pustaka Setia
- Bassola, M., Tazza, L., Pannocchia, N., Ciciarelli, A., Liberatoru, M., & Luciani, G. (2001), Malnutrition in Hemodialysis Patients. Diunduh tanggal 14 Desember 2015. <http://ncbi.nlm.nih.gov>
- Black, J.M., & Hawks, J.H. (2009). *Medical Surgical Nursing Clinical Management or Positive Outcome*. (8th ed). St. Louis: Elsevier
- Bridge, A. B., & Holt, K. E. (2015). Effect of nocturnal dialysis on prognosis in adult patients: A review of the literature. *Nephrology Nursing Journal*, 42(4), 375-380. Retrieved from <http://search.proquest.com/docview/1705656159?accountid=38628>
- Broschius, S.K & Castagnola J. (2006). Chronic Kidney Disease: acute manifestations and role of critical care nurses. *Critical Care Nurse*.

- Aug.26 (4): 17-20, 22-7. Diunduh tanggal 14 Desember 2015.
<http://ncbi.nlm.nih.gov>
- Bonner, A., Wellard, S., & Caltabiano, M. (2010). The impact of fatigue on daily activity in people with chronic kidney disease. *Journal of Clinical Nursing*, 19(21-22), 3006-3015. Diunduh tanggal 15 November 2016
<http://search.proquest.com/docview/1705656159?accountid=38628>
- Brunner, L.S. & Suddarth, D.S. (2010). *Buku Ajar Keperawatan Medikal Bedah*. Edisi 8. Jakarta: Kedokteran ECG.
- Chang, Y., Cheng, S., Lin, M., Gau, F., & Chao, Y. C. (2010). The effectiveness of intradialytic leg ergometry exercise for improving sedentary life style and fatigue among patients with chronic kidney disease: A randomized clinical trial. *International Journal Of Nursing Studies*, 47(11), 1383-1388. doi:10.1016/j.ijnurstu.2010.05.002
- Cheena, B. S., O'Sullivan. J, Chan, M., Patwardhan A., *et.al.* (2006). *Progressive resistance training during hemodialysis : rationale and method of a randomized-controlled trial. Hemodialysis International Journal*; 10: 303-310. Diunduh tanggal 29 Juni 2016 dari <http://www.interscience.com>
- Cupisti, A., D'alessandro, C., Bottai, A., Fumagalli, G., & Capitanini, A. (2013). Physical activity and exercise training: A relevant aspect of the dialysis patient's care. *Internal and Emergency Medicine*, 8, 31-4. doi:<http://dx.doi.org/10.1007/s11739-013-0917-y>
- Corwin, E.J. (2009). *Buku Saku Patofisiologi*. Jakarta : EGC
- Danismaya, Irawan, 2008. Pengaruh Tehnik Relaksasi Yoga terhadap *Fatigue* penderita kanker pasca kemoterapi di RS Hasan Sadikin Bandung, (tesis). Perpustakaan UI
- Daniyati, P. (2015). Evidence Based Practice Intradialytic Exercise Untuk Pengelolaan Tekanan Darah Pada Pasien Chronic Kidney Disease (Ckd) Stage V., diakses tanggal 16 Maret 2017. jurnal.akper-notokusumo.ac.id/index.php/jkn
- Danquah, F., Zimmerman, L., Diamond, P.M., Meininger, J., & Bergstrom, N. (2010). Frequency, severity, and distress of dialysis-related

symptoms reported by patients on hemodialysis. *Nephrology Nursing Journal*, 37(6), 627-638.

- Daugirdas, J.T., Blake, P.G. (2007). *Handbook of Dialysis*. (4th ed). Lippincot:Philadelphia
- De-Lima, M.C., Cicotoste, C., Cardoso, K.D., Junior, L.G.F., Monteiro, M.B., & Dias AS. (2013). Effect of exercise performed during hemodialysis: Strength versus Aerobic. *Renal Failure*, 35(5), 697–704. 2800.20130028.
- Da-Silva, S.F., Pereira, A.A., Aparecido, W., Silva, H., d., Simões, R., Barros, J., d. R., & Neto. (2013). Physical therapy during hemodialysis in patients with chronic kidney disease. *J Bras Nefrol*, 35. doi: 10.5935/0101
- Evans, W.J., & Lambert, C.P. (2007). Physiological basis of fatigue. *American Journal of Physical Medicine & Rehabilitation*, 86(1, Suppl.), S29-S46. Retrieved from <http://search.proquest.com/docview/1041232712?accountid=38628>
- Firmansyah, A.M. (2010). Usaha memperlambat perburukan penyakit ginjal kronik ke penyakit ginjal stadium akhir. *Cermin Dunia Kedokteran*; ISSN: 0125-913 X/176/vol 37 No 3/April 2010
- Fritz S., (2005). *Sport & Exercise Massage*, St., Louis Missouri: Elsevier Mosby
- Guyton, A. & Hall, J. (2007). *Buku ajar fisiologi kedokteran*. Jakarta : EGC
- Girija, K & Radha, R. 2013. Beneficial Effect of Physical Activity in Hemodialysis Patients. *Universal Journal of Engineering Science* .Retrieved from: <http://www.hrpub.org>
- Henson, A., Gillespie, B., McCarthy, A., Finch, L., Chatterton, S., Devlin, J., .Orazio, L. (2010). Intradialytic exercise: A feasibility study. *Renal Society of Australasia Journal*, 6(1), 11-15.
- Horigan, A. E. (2012). The experience and self-management of fatigue in adult hemodialysis patients. (Order No. 3523263, Duke University).

ProQuest Dissertations and Theses, 165-n/a. Retrieved from <http://search.proquest.com/docview/1041232712?accountid=38628>

- Horigan, A., Rocchiccioli, J., & Trimm, D. (2012). Dialysis and fatigue: Implications for nurses - A case study analysis. *Medsurg Nursing*, 21(3), 158-63, 175. Retrieved from <http://search.proquest.com/docview/1021184805?accountid=38628>
- Ignatavicius. D.D, & Workman. L.M., (2006). *Medical surgical nursing critical thinking for collaboration care* (5th).St.Louis.Elsevier
- Jablonski, A. (2007). The multidimensional characteristics of symptoms reported by patient on hemodialysis. *Nephrology Nursing Journal*.34 (1).29
- Johansen. (2005). *Exercise and chronic kidney disease : current recommendations*.*Sports Med* 2005;35(6):485-99. Diunduh tanggal 10 Desember 2015, dari www.jasn.org
- Jung TD, Park SH. (2011). *Intradialytic exercise programs for hemodialysis patients*. *Chonnam Med J*. 2011;**47**(2):61–5. Diunduh tanggal 30 Januari 2016, dari <http://www.proquest.umi.com>
- Kementerian Kesehatan Republik Indonesia.(2011). Pedoman interpretasi data klinik.
- Knap B, Ponikvar B.J, Ponikvar R, Bren F.A. (2005). *Regular exercise as a part of treatment for patients with end stage renal disease*. *Therapeutic Apheresis and Dialysis*; 9 (3):211-213, Diunduh tanggal 30 Januari 2016, dari <http://www.proquest.umi.com>
- Kring, D.L & Crane. (2009). Factors affecting Quality of life in persons on hemodialysis, *Nephrology Nursing Journal*, 36, 15-55. Diunduh tanggal 5 Desember 2015, dari <http://www.proquest.umi.com>
- Kusmana. (2007). *Olah raga untuk orang sehat dan penderita penyakit jantung*.FKUI: Jakarta
- Lee, G. R. (2000). The Normocytic Normochromic Anemias dalam G. R. Lee, T. C. Bithel, J. Foeter, J. Athens, J. N. Lukens (eds): *Wintrobe Clinical Hematology 9th edition*. Vol. 1. Lea & Febriger. London

- Locatelli, F., Fauque, D., Heimbürger O. (2002). Nutritional Status in dialysis patients: European consensus. *Nephrology dialysis Transplantation*, 17, 563-572
- Maheshwari, V., Samavedham, L., Rangaiyah, Gade P., Loy, Y., Ling, Lieng H., Sethi, S., Leong Titus L. W. (2012). Comparison of toxin removal outcomes in online hemodiafiltration and intradialytic exercise in high-flux hemodialysis : A prospective randomized open-label clinical study protocol. *BMC Nephrology* 13:156. doi:10.1186/1471-2369-13-156.
- Makhlough, A., Ilali, E., Mohseni, R., & Shahmohammadi, S. (2012). Effect of Intradialytic Aerobic Exercise on Serum Electrolytes Levels in Hemodialysis Patients. *Iranian Journal Of Kidney Diseases*, 6(2), 119-123.
- Maulidya, Nindy. (2015). Gambaran Jenis Anemia Menggunakan Mean Corpuscular Hemoglobin (MCH) pada Gagal Ginjal Kronik.
- McCann, K., & Heuter, J.R. (2010). Fatigue in persons with renal failure who require maintenance haemodialysis. *Journal of Advanced Nursing*, 32(5),1132-1142. Diunduh tanggal 5 Desember 2015, dari <http://www.proquest.umi.com>
- Mitchell, S.A., Beck, S.L., Hood, L., Moore, K., Tanmer, E.R. (2007). *Putting evidence into practice : Evidence based intervention for fatigue during and following cancer and its treatment. Clinical Journal of Oncology Nursing*, 11 (1), 99-113
- Mohseni, R., Zeydi, A. E., Ilali, E., Adib-Hajbaghery, M., & Makhlough, A. (2013). The Effect of Intradialytic Aerobic Exercise on Dialysis Efficacy in Hemodialysis Patients: A Randomized Controlled Trial. *Oman Medical Journal*, 28(5), 345-349. doi:10. 5001/omj.2013.99
- Mortazavi, M., Vahdatpour, B., Ghasempour, A., Taheri, D., Shahidi, S., & Moeinzadeh, F. (2013). Aerobic exercise improves signs of restless leg syndrome in end stage renal disease patients suffering chronic hemodialysis. *The Scientific World Journal*, 1-4.
- Murray RK, Granner DK, Rodwell VW. (2009). *Biokimia Harper*. Jakarta: Penerbit Buku Kedokteran EGC

- Nasution, R.S. (2010). Latihan jasmani pada pasien hemodialisis regular. Diunduh pada tanggal 23 Januari 2016 dari <http://nefrologyners./latihan-pada-pasien-hemodialisis-reguler>.
- National Kidney Foundation (NKF) Kidney Disease Outcome Quality Initiative (K/DOQI)*, (2002). Advisory board: K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification and stratification. *Kidney disease outcome quality initiative*. *Am J Kidney Dis* 2005;39(Suppl 1):S1-246.
- Nekada, Cornelia. D.Y. (2014). Pengaruh Gabungan Relaksasi Napas Dalam dan Otot Progresif terhadap Komplikasi Intradialisis di unit hemodialisis RSUP Dr. Soeradji Tirtonegoro klaten. Tesis.Unpad
- Nurchayati, Sofiana. (2010). Faktor-Faktor Yang Berhubungan Dengan Kualitas Hidup Pasien Penyakit Ginjal Kronik Yang Menjalani Hemodialisis di Rumah Sakit Umum Daerah Banyumas. Tesis: UI
- O’Sullivan, D., & McCarthy, G. (2009). *Exploring the symptom of fatigue inpatients with end stage renal disease*. *Nephrology Nursing Journal*. 37-47. Diunduh tanggal 30 Januari 2016, dari <http://www.proquest.umi.com>
- Orti. E.S., (2010). Exercise in Hemodialysis patients : A literature Systematic review. *Nefrologi*. diunduh tanggal 1 Januari 2016. diperoleh dari <http://revistanefrologia.com>
- Painter, P., Moore, G., Carlson, L., Paul, S., Myll, J., Phillips, W., & Haskell, W. (2000). Exercise A Guide for the People on Dialysis. *American Journal of Kidney Diseases*, Diunduh tanggal 15 November 2016. www.proquest.umi.com
- Pali, D.V. (2012). Gambaran anemia pada penderita penyakit ginjal kronik di blu. Rsup. Prof. Dr. R. D. Kandou. Available from: <http://ejournal.unsrat.ac.id/index.php/eclinic/article/view/3282>
- Parker, K., Zhang, X., Lewin, A., & MacRae, J. M. (2015). The association between intradialytic exercise and hospital usage among hemodialysis patients. *Applied Physiology, Nutrition & Metabolism*, 40(4), 371-378. doi:10.1139/apnm-2014-0326

- Parsons, T.K., Tosselmire E.D., King-VanVlack C.E. (2006). *Exercise training during hemodialysis improves dialysis efficacy and physical performance. Exercise Arch phys med Rehabil*: 2006; 87:680-7, from <http://www.interscience.com>
- PERNEFRI. (2003). *Konsensus Dialisis Perhimpunan Nefrologi Indonesia*. Jakarta
- PERNEFRI. (2012). *Naskah Lengkap Workshop Dan Simposium Nasional Peningkatan Pelayanan Hemodialisis, Penyakit Ginjal Dan Aplikasi Indonesian Renal Registry*
- Piper BF, Dibble SL, Dodd MJ, Weiss MC, Slaughter RE, Paul SM. (1998) The revised Piper Fatigue Scale: psychometric evaluation in women with breast cancer. *Oncol Nurs Forum*.
- Potter & Perry (2006). *Renata Kumalasari dkk, (alih bahasa) Fundamental Keperawatan*: EGC: Jakarta
- Price. (2006). *Patofisiologi konsep klinis proses-proses penyakit*. EGC.Jakarta
- Pujiastuti, T., (2014). *Pengaruh Exercise Intradialisis Terhadap Peningkatan Adekuasi Hemodialisis Pasien Chronic Kidney Disease Di Rumah Sakit Panti Rapih Yogyakarta*
- Rahardjo P, Susalit E, Suhardjono. (2001). *Dialisis dalam Buku Ajar Ilmu Penyakit Dalam (Jilid II)*.Balai Penerbit FKUI.Jakarta
- Rosenthal M. (2005). *The complete thyroid book*, Mc Graw Hill, New York
- Ross, A.F & Kearney. (2000). *Dialysis Disequilibrium Syndrome. American Journal of Nursing*. 100, 53-54.
- Runtung Y, Kadir A, dan Akuilina S.(2013). *Pengaruh hemodialisa terhadap kadar ureum kreatinin dan hemoglobin pada pasien GGK di Ruang Haemodialisa RSUP DR Wahidin Sudirohusodo Makassar*;2(3):1-7.
- Sastroasmoro, S. dan Ismael, S. (2011). *Dasar-dasar Metodologi Penelitian Klinis*.Edisi ke-3.Jakarta: Sagung Seto
- Sheng, K., Zhang, P., Chen, L., Cheng, J., Wu, C., & Chen, J. (2014). *Intradialytic exercise in hemodialysis patients: A systematic review*

- and meta-analysis. *American Journal of Nephrology*, 40(5), 478-490. doi:<http://dx.doi.org/10.1159/000368722>
- Smeltzer, S.C., & Bare. B.G., (2008). *Textbook Of Medical Surgical Nursing*. (1st Ed). Philladelphia : Lipincott
- Solimun, M. H. (2015). Effect of intradialytic exercise on fatigue, electrolytes level and blood pressure in hemodialysis patients: A randomized controlled trial. *Journal of Nursing Education and Practice*, 5(11), 16. doi:<http://dx.doi.org/10.5430/jnep.v5n11p16>
- Sudoyo, A.W., Setiyohadi, B., Alwi I., Simadibrata, M., & Setiati, S. (2009). *Buku Ajar Ilmu Penyakit Dalam*. (Edisi V). Jilid II. Pusat Penerbitan Ilmu Penyakit Dalam FKUI : Jakarta
- Sulisyarningsih, R, D. (2010). Efektivitas latihan fisik selama hemodialisis terhadap peningkatan kekuatan otot pasien penyakit ginjal kronik di Rumah Sakit Umum Daerah Kota Semarang.
- Sullivan, D & McCarthy, G. (2009). Exploring the symptom of fatigue in patients with end stage renal disease. *Nephrology Nursing Journal*. 36, 38-40. Renal Nursing. 2th edition. Philadelphia : Elsevier Science
- Supranto, J. (2013). *Teknik Sampling untuk Survei dan Eksperimen*. Penerbit PT Rineka Cipta, Jakarta.
- Takhreem, M., (2008) *The effectiveness of intradialytic exercise prescription on quality of life in patient with chronic kidney disease*. *Medscape J Med*. 2008;10 (10): 228, diunduh tanggal 27 Januari 2016 <http://ncbi.nlm.nih.gov>
- Thomas-Hawkins, C., & Zazworsky, D. (2003). Self-management of chronic kidney disease: Patients shoulder the responsibility for day-to-day management of chronic illness. How can nurses support their autonomy? . *American Journal of Nursing*, 10 5(10), 40- 48.
- Weisbord SD, Fried LF, Mor MK *et al*. Renal provider recognition of symptoms in patients on maintenance hemodialysis. *Clin. J. Am. Soc. Nephrol*. 2007; 2: 960–967.
- Wong, S.W, Chan, Y.M,& Lim T.S. (2011). Correlates of Physical Activity Level among Hemodialysis Patients in Selangor, Malaysia. Malaysia: Universiti Putra Malaysia.