

DAFTAR PUSTAKA

- Aminullah, A., & Gatot, D. (2007). *Penatalaksanaan Sepsis Neonatorum*. Jakarta: Departemen Kesehatan Republik Indonesia. 23 Mei 2016.
- Baratawidjaja, K. G., & Rengganis, I. (2014). *Imunologi Dasar* (11th ed.). Jakrta: Badan Penerbit FKUI.
- Behrman, R. E., Kliegman, R., & Arvin, A. M. (2000). *Nelson Textbook of Pediatrics*. (A. S. Wahab, Noerhayati, H. Soebono, Sonarto, Sunartini, M. Juffrie, M. Julia, Eds.) (15th ed.). Philadelphia: W. B. Saunders Company.
- Bhandari, V., Wang, C., Rinder, C., & Rinder, H. (2008). Hematologic Profile of Sepsis in Neonates : Neutrophil CD64 as a Diagnostic Marker, 121(1). <http://doi.org/10.1542/peds.2007-1308>. Diakses pada: 30 April 2016.
- Carolus, W., Rompis, J., & Wilar, R. (2013). HUBUNGAN APGAR SKOR DAN BERAT BADAN LAHIR DENGAN SEPSIS NEONATORUM, 1-7.<https://ejournal.unsrat.ac.id/index.php/eclinic/article/view/3271/2815>Diakses pada:13 Mei 2017.
- Christensen, A. R. D., & Baer, V. L. (2012). Reference Ranges for Lymphocyte Counts of Neonates : Associations Between Abnormal Counts and Outcomes, 129(5). <http://doi.org/10.1542/peds.2011-2661>. Diakses pada: 23 Mei 2016.
- Darnifayanti, Tjipta, G. D., Rusdijas, & Lubis, B. M. (2009). Immature-to-total neutrophil ratio as an early diagnostic tool of bacterial neonatal sepsis. *Pediatrica Indonesiana*, 49(2), 97–103. Retrieved from <http://paediatricaindonesiana.org/pdffile/55-3-7.pdf>. Diakses pada: 23 Mei 2016.
- Dorland, W. A. N. (2007). *Dorland's Illustrated Medical Dictionary* (31st ed.). Singapore: Elsevier.
- Enguix, A., Rey, C., Concha, A., Medina, A., Coto, D., & Dieguez, M. A. (2000). Comparison of procalcitonin with C-reactive protein and serum amyloid for the early diagnosis of bacterial sepsis in critically ill neonates and children, (April 2016). <http://doi.org/10.1007/s001340000709>. Diakses pada: 30 April 2016.
- Fraser, T., Tilyard, M., Colquhoun, D., Ikram, R., Jensen, P., Kyle, C., ... Reith, D. (2008). Complete Blood Count. *Bpac (Nz)*, 1–24. Retrieved from http://www.bpac.org.nz/Supplement/2008/May/docs/bpac_cbc_in_primary_care.pdf. Diakses pada: 16 Juni 2016.
- Grope, I., & Gardovska, D. (2009). Diagnostic Markers for Early Sepsis Diagnosis in Children with Systemic Inflammatory Response Syndrome, 63(4), 191–197. <http://doi.org/10.2478/v10046-009-0048-z>. Diakses pada: 21 Mei 2016.
- Guyton, A. C., & Hall, J. E. (2014). *Textbook of Medical Physiolgy* (11th ed.).
- Holcomb, I. N., Kabakoff, R. C., Chan, B., Baker, T. W., Gurney, A., Henzel, W., ... Hébert, C. C. (2000). FIZZ1, a novel cysteine-rich secreted protein associated with pulmonary inflammation, defines a new gene family. *The EMBO Journal*,

19(15), 4046–55. <http://doi.org/10.1093/emboj/19.15.4046>. Diakses pada: 23 Mei 2016.

Hotchkiss, R. S., & Karl, I. E. (2003). The Pathophysiology and Treatment of Sepsis, 138–150. Retrieved from <http://www.nejm.org/doi/full/10.1056/NEJMra021333>. Diakses pada: 23 Mei 2016.

Jakubik, L. D. (2014). Sharpen Your Pediatric Laboratory Interpretation Skills: Pediatric Complete Blood Count (CBC), Differential, and Absolute Neutrophil Count (ANC). *Pediatric Nursing Annual Conference*. Retrieved from http://www.pediatricnursing.net/conf/handouts/2010/session_121_jakubik.pdf. Diakses pada: 16 Maret 2016.

Klingenberg, C. (2005). Coagulase-negative staphylococcal sepsis in neonates : Association between antibiotic resistance , biofilm formation and the host Coagulase-Negative Staphylococcal Sepsis in Neonates, (April 2016). <http://doi.org/10.1097/01.inf.0000176735.20008.cd>. Diakses pada: 30 April 2016.

Kocherlakota, P., & Gamma, E. F. La. (1997). Human Granulocyte Colony-stimulating Factor May Improve Outcome Attributable to Neonatal Sepsis Complicated by Neutropenia, 100(1). Retrieved from <http://pediatrics.aappublications.org/content/100/1/e6>. Diakses pada: 5 Mei 2016.

Kurt, A., Aygun, A., Godekmerdan, A., Kurt, A., Dogan, Y., & Yilmaz, E. (2007). Serum IL-1 beta, IL-6, IL-8, and TNF-alpha levels in early diagnosis and management of neonatal sepsis. *Mediators of Inflammation*, 2007, 1–5. <http://doi.org/10.1155/2007/31397>. Diakses pada: 16 Maret 2016.

Lockwood, W. (2015). The Complete Blood Count and Associated Tests Course objectives. Retrieved from <http://www.rn.org/courses/coursematerial-263.pdf>. Diakses pada: 5 Mei 2016.

Mayuga, W. A. B., & Isleta, P. F. D. (2005). Clinical Correlation of Neonatal and Maternal Hematological Parameters as Predictors of Neonatal Sepsis, 9(2), 36–43. Retrieved from https://www.researchgate.net/publication/267554015_CLINICAL_CORRELATION_OF_NEONATAL_AND_MATERNAL_HEMATOLOGICAL_PARAMETERS_AS_PREDICTORS_OF_NEONATAL_SEPSIS. Diakses pada: 23 Mei 2016.

Mehta, A., & Hoffbrand, V. (2006). *At a Glance: Hematologi*. Jakarta: EMS.

Murphy, K. (2011). *Janeway's Immunobiology* (8th ed.). New York: Garland Science.

Newman, T. B., Puopolo, K. M., Wi, S., Draper, D., & Escobar, G. J. (2010). Interpreting complete blood counts soon after birth in newborns at risk for sepsis.

Pediatrics, 126(5), 903–909. <http://doi.org/10.1542/peds.2010-0935>. Diakses pada: 5 Mei 2016.

Pablo, R. De, Monserrat, J., Prieto, A., & Alvarez-mon, M. (2014). Role of Circulating Lymphocytes in Patients with Sepsis, 2014. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/25302303>. Diakses pada: 21 Mei 2016.

Paolucci, M., Corvaglia, L., & Landini, M. P. (2016). Laboratory diagnosis of late-onset sepsis in newborns by multiplex real-time PCR, 533–534. <http://doi.org/10.1099/jmm.0.003848-0>. Diakses pada: 30 April 2016.

Punt, J., Owen, J., & Caligiuri, M. A. (2001). The biology of human natural killer-cell subsets. *Trends in Immunology*, 22(11), 633–640. [http://doi.org/10.1016/S1471-4906\(01\)02060-9](http://doi.org/10.1016/S1471-4906(01)02060-9). Diakses pada: 11 Juni 2016.

Pusponegoro, T. S. (2000). Sepsis pada Neonatus (Sepsis Neonatal). *Sari Pediatri*, 2, 96–102. <https://saripediatri.org/index.php/saripediatri/article/download/1038/968> . Diakses pada: 13 Mei 2017

Putra, P. J. (2012). Insiden dan Faktor-Faktor yang Berhubungan dengan Sepsis Neonatus di RSUP Sanglah Denpasar. *Sari Pediatri*, 14(3), 205–210. Retrieved from <http://saripediatri.idai.or.id/pdf/14-3-12.pdf>. Diakses pada: 30 April 2016.

Roeslani, R. D., Amir, I., & Nasrulloh, M. H. (2013). Penelitian Awal: Faktor Risiko pada Sepsis Neonatorum Awitan Dini. *Sari Pediatri*, 14(6), 363–368. Retrieved from <http://saripediatri.idai.or.id/pdf/14-6-6.pdf>. Diakses pada: 30 April 2016.

Shah, B. A., & Padbury, J. F. (2014). Neonatal sepsis: an old problem with new insights. *Virulence*, 5(1), 170–8. <http://doi.org/10.4161/viru.26906>. Diakses pada: 26 Mei 2016

Simbolon, D. (2008). Faktor risiko sepsis pada bayi baru lahir di RSUD Curup kabupaten Rejang Lebong. *Buletin Penelitian Kesehatan*, (1 994). Retrieved from <http://ejournal.litbang.depkes.go.id/index.php/BPK/article/download/2166/1118>. Diakses pada: 23 Mei 2016.

Sundaram, V., Kumar, P., Dutta, S., Mukhopadhyay, K., Ray, P., Gautam, V., & Narang, A. (2009). Blood Culture Confirmed Bacterial Sepsis in Neonates in a North Indian Tertiary Care Center: Changes over the Last Decade, 46–50. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19168958>. Diakses pada: 30 April 2016.

Suryaatmadja, M., Augustine, M., Salim, K. A., Tobing, D. L., Gunadi, D., Haryanto, T., ... Hwa, T. S. (2004). Tabel Konversi Sistem Satuan SI - Konvensional dan Nilai Rujukan Dewasa - Anak Parameter Laboratorium Klinik. In M. Suryaatmadja (Ed.). Jakarta: Perhimpunan Dokter Spesialis Patologi Klinik Indonesia.

Utomo, M. T. (2010). Risk Factors of Neonatal Sepsis : A Preliminary Study in. *Indonesian Journal of Tropical and Infectious Disease*, 1(1). <http://ejournal.unair.ac.id/index.php/IJTID/article/download/3718/2522>. Diakses pada:23 Mei 2016

Weinberg, A. G., Rosenfeld, C. R., Manroe, B. L., & Browne, R. (1985). Neonatal blood cell count in health and disease. II. Values for lymphocytes, monocytes, and eosinophils. *The Journal of Pediatrics*, 106(3), 462–466. [http://doi.org/10.1016/S0022-3476\(85\)80681-8](http://doi.org/10.1016/S0022-3476(85)80681-8). Diakses pada: 23 Mei 2016.

Yasa, I. W. P. S. (2014). biomarker pada Sepsis Neonatal. *Symposium SURAMADE, Patologi Klinik*.

Zahedpasha, Y., Kacho, M. A., & Hajiahmadi, M. (2009). Procalcitonin as a Marker of Neonatal Sepsis, 19(2), 117–122. Retrieved from <https://tspace.library.utoronto.ca/bitstream/1807/58849/1/pe09018.pdf>. Diakses pada: 30 April 2016.

Zakaria, M., & Rafaat, M. (2012). Immunophenotyping of Lymphocyte Subpopulations and pre-inflammatory mediators in neonatal sepsis. *Advances in Environmental Biology*, 6(11), 2948–2952. Retrieved from <http://www.aensiweb.com/old/aeb/2012/2948-2952.pdf> . Diakses pada: 23 Mei 2016.