

DAFTAR PUSTAKA

- Al-azzawi, S. I., Prof, A. and Salal, R. H. (2013) ‘Influence of age and gender on salivary flow rate in completely edentulous patients’, *MDJ*, Vol.:10 No(1), pp. 64–68.
- Alam, M. K. (2014) *A to Z Orthodontics. Volume 3: Malocclusion*.
- Alves, C., Brandão, M., Andion, J. and Menezes, R. (2010) ‘Use of graduated syringes for measuring salivary flow rate: A pilot study’, *Brazilian Dental Journal*, 21(5), pp. 401–404. doi: 10.1590/S0103-64402010000500004.
- AR, Prabhakar; Gulati, Akanksha; Mehta, Deepak; S, S. (2009) ‘Diagnostic Applications of Saliva in Dentistry 1 1’, *International Journal of Clinical Pediatric Dentistry*, 2(December), pp. 7–13.
- Arab, S., Malekshah, S. N., Mehrizi, E. A. and Khanghah, A. E. (2016) ‘Effect of Fixed Orthodontic Treatment on Salivary Flow , pH and Microbial Count’, 13(1).
- Bahreman, A. (1972) ‘Orthodontic treatment’, *Zahnärztliche Mitteilungen*, 62(18), pp. 903–907.
- Benson, P. E., Javidi, H. and DiBiase, A. T. (2015) ‘What is the value of orthodontic treatment?’, *BDJ*, 218(3), pp. 185–190. doi: 10.1038/sj.bdj.2015.43.
- Berti-Couto, S. de A., Couto-Souza, P. H., Jacobs, R., Nackaerts, O., Rubira-Bullen, I. R. F., Westphalen, F. H., Moysés, S. J., Ignácio, S. A., Costa, M. B. da and Tolazzi, A. L. (2012) ‘Clinical diagnosis of hyposalivation in hospitalized patients.’, *Journal of applied oral science : revista FOB*, 20(2), pp. 157–61. doi: 10.1590/S1678-77572012000200006.
- Bonetti, G. A., Parenti, S. I., Garulli, G., Gatto, M. R. and Checchi, L. (2013) ‘Effect of fixed orthodontic appliances on salivary properties’, pp. 12–15.
- Carvalho, P. M., Castelo, P. M., Carpenter, G. H. and Gavião, M. B. D. (2016) ‘Masticatory function , taste , and salivary flow in young healthy adults’, 58(3), pp. 391–399. doi: 10.2334/josnusd.16-0135.

- Cheng, L. L., Moor, S. L., Kravchuk, O., Meyers, I. A. and Ho, C. T. C. (2007) ‘Bacteria and salivary profile of adolescents with and without cleft lip and/or palate undergoing orthodontic treatment’, *Australian Dental Journal*, 52(4), pp. 315–321. doi: 10.1111/j.1834-7819.2007.tb00508.x.
- Cobourne, M.T. & DiBiase, A.T., 2010. *Handbook of Orthodontics*. China: Mosby Elsevier.
- Dawes, C. (1972) ‘Circadian rhythms in human salivary flow rate and composition’, *J Physiol*, 220(3), pp. 529–545.
- De Almeida, P. D. V., Grégio, A. M. T., Machado, M. Â. N., De Lima, A. A. S. and Azevedo, L. R. (2008) ‘Saliva composition and functions: A comprehensive review’, *Journal of Contemporary Dental Practice*, 9(3), pp. 072–080. doi: 1526-3711-497 [pii].
- Farnaud, S. J. C., Kosti, O., Getting, S. J. and Renshaw, D. (2010) ‘Saliva: Physiology and Diagnostic Potential in Health and Disease’, pp. 434–456. doi: 10.1100/tsw.2010.38.
- Foster, T.D., 1997. Buku Ajar Ortodonsi. Translated by L. Yuwono. Jakarta: EGC.
- Hall, J. E. and Guyton, A. C. (2011) *Guyton and Hall Textbook of Medical Physiology, Journal of Chemical Information and Modeling*. doi: 10.1017/CBO9781107415324.004.
- Humphrey, S. P. and Williamson, R. T. (2001) ‘A review of saliva: Normal composition, flow, and function’, *Journal of Prosthetic Dentistry*, 85(2), pp. 162–169. doi: 10.1067/mpr.2001.113778.
- Indriana, T., Kalimantan, J., Kampus, N. and Boto, T. (2011) ‘Perbedaan Laju Aliran Saliva dan pH karena Pengaruh Stimulus Kimiaawi dan Mekanis’, 17(44), pp. 1–5.
- Isaacson, K.G., Muir, J.D. & Reed, R.T., 2007. *Removable Orthodontic Appliances*. New Delhi, India: Elshiever.
- Joskow, J. and Yamane, T. (1965) ‘Statistics, an Introductory Analysis.’, *Journal of the American Statistical Association*, 60(310), p. 678. doi: 10.2307/2282703.
- Karnik, A., Pagare, S., Krishnamurthy, V., Vahanwala, S. and Waghmare, M.

- (2015) ‘Determination of salivary flow rate, pH, and dental caries during pregnancy: A study’, *Journal of Indian Academy of Oral Medicine and Radiology*, 27(3), p. 372. doi: 10.4103/0972-1363.170454.
- Kasuma, N., 2015. Fisiologi dan Patologi Saliva. Padang: Andalas University Press. p.25.
- Kidd, E.A.M. & Joyston-Bechal, S., 2013. *Dasar - Dasar Karies Penyakit dan Penanggulangan*. Translated by N. Sumawinata & S. Faruk. Jakarta: EGC.
- Kundu, R., Tripathi, A. M., Jaiswal, J. N., Ghoshal, U. and Palit, M. (2016) ‘Effect of fixed space maintainers and removable appliances on oral microflora in children : An in vivo study’, pp. 3–9. doi: 10.4103/0970-4388.175498.
- Lara-carrillo, E., Sánchez-pérez, L. and Alanís-tavira, J. (2010) ‘Effect of orthodontic treatment on saliva , plaque and the levels of Streptococcus mutans and Lactobacillus’, 15(6). doi: 10.4317/medoral.15.e924.
- Mitchell, L. (2013) ‘The Rationale for Orthodontic Treatment. Introduction to Orthodontics.’, *Journal of Chemical Information and Modeling*, pp. 2–8. doi: 10.1017/CBO9781107415324.004.
- Mitchell, L., Mitchell, D.A. & McCaul, L., 2014. *Kedokteran Gigi Klinik*. 5th ed. Jakarta: EGC.
- Navazesh, M. and Kumar, S. K. S. (2008) ‘Measuring salivary flow: challenges and opportunities.’, *Journal of the American Dental Association* (1939). © 2008 American Dental Association, 139 Suppl(May), p. 35S–40S. doi: 10.14219/jada.archive.2008.0353.
- Peros, K., Mestrovic, S., Anic-Milosevic, S. and Slaj, M. (2011) ‘Salivary microbial and nonmicrobial parameters in children with fixed orthodontic appliances’, *The Angle Orthodontist*. Edward H. Angle Society of Orthodontists, 81(5), pp. 901–906. doi: 10.2319/012111-44.1.
- Preoteasa, C. T., Ionescu, E. and Preoteasa, E. (2012) ‘Risks and Complications Associated with Orthodontic Treatment’, *Orthodontics – Basic Aspects and Clinical Considerations*, 18, pp. 403–428.
- Saloom, H. F., Mohammed-Salih, H. S. and Rasheed, S. F. (2013) ‘The influence

- of different types of fixed orthodontic appliance on the growth and adherence of microorganisms (in vitro study)', *Journal of Clinical and Experimental Dentistry*, 5(1), pp. 36–41. doi: 10.4317/jced.50988.
- Sánchez, E. R. B. and Honores, M. J. C. (2015) 'Effect of orthodontic fixed appliances on salivary flow and viscosity', 3, pp. 186–190. doi: 10.1016/j.rmo.2016.03.049.
- Shetty, C., Hegde, M. N. and Devadiga, D. (2013) 'Correlation between dental caries with salivary flow, pH, and buffering capacity in adult south Indian population: An in-vivo study', *International Journal of Research in Ayurveda and Pharmacy*, 4(2), pp. 219–223. doi: 10.7897/2277-4343.04226.
- Ship, J. A. and Fischer, D. J. (1997) 'The relationship between dehydration and parotid salivary gland function in young and older healthy adults', *Journals of Gerontology Series A-Biological Sciences & Medical Sciences*, 52(5), pp. M310–M319. Available at: 9310086.
- Shukla, C., Maurya, R. K., Singh, V. and Tijare, M. (2016) 'Evaluation of changes in Streptococcus mutans colonies in microflora of the Indian population with fixed orthodontics appliances', pp. 309–314.
- Singh, G. (2007) *Textbook of orthodontics*. 2nd edn, Jaypee, India. 2nd edn. doi: 10.1017/CBO9781107415324.004.
- Takeuchi, K., Furuta, M., Takeshita, T., Shibata, Y., Shimazaki, Y., Akifusa, S., Ninomiya, T., Kiyohara, Y. and Yamashita, Y. (2015) 'Risk factors for reduced salivary flow rate in a Japanese population: The Hisayama study', *BioMed Research International*, 2015. doi: 10.1155/2015/381821.
- Waugh, A. and Grant, A. (Allison W. (2014) *Ross and Wilson anatomy & physiology in health and illness*, Churchill Livingston Elsevier. doi: 10.1017/CBO9781107415324.004.