

DAFTAR PUSTAKA

- Ahmed, S. E. S., & Younis, G. a. (2014). The effect of Relaxation Techniques on Quality of Sleep for Patients with End-Stage Renal Failure Undergoing Hemodialysis. *International Journal of Innovative And Applied Research*, 2(7), 1–12. diakses 01 Oktober 2017 dari http://www.journalijiar.com/uploads/2014-07-31_102535_59.pdf
- Akgun, Z. N. & Dayapoglu, N. (2015). Effect of progressive relaxation exercises on fatigue and sleep quality in patients with chronic obstructive lung disease (COPD). *Complementary Therapies in Clinical Practice*, 21(4), 277–281. <https://doi.org/10.1016/j.ctcp.2015.10.002> diakses 17 Juli 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/26573455>
- Akyol, A. D., Yildirim, Y., Toker, E., & Yavuz, B. (2011). The use of complementary and alternative medicine among chronic renal failure patients. *Journal of Clinical Nursing*, 20(7–8), 1035–1043. <https://doi.org/10.1111/j.1365-2702.2010.03498.x> diakses 10 Oktober 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/21320219>
- Alligood, MR & Tomey AN. (2014).*Nursing Theorist and Their Work, Sixth Edition*, St. Louis: Mosby Elsevier, Inc
- Amini, E., Goudarzi, I., Masoudi, R., Ahmadi, A., & Momeni, A. (2016). Effect of Progressive Muscle Relaxation and Aerobic Exercise on Anxiety, Sleep Quality, and Fatigue in Patients with Chronic Renal Failure Undergoing Hemodialysis, 8(12), 1634–1639. diakses 10 Juli 2017 dari http://impactfactor.org/PDF/IJPCR/8/IJPCR_Vol8_Issue12_Article14.pdf

Arif Muttaqin, K. S. (2011). *Asuhan keperawatan gangguan sistem perkemihan*. Jakarta: Salemba Medika

Arifin, Zainal. (2011). *Analisis hubungan kualitas tidur dengan kadar glukosa darah pasien diabetes mellitus tipe 2 di rumah sakit umum propinsi Nusa Tenggara Barat*. Tesis. Universitas Indonesia

Badan Penelitian dan Pengembangan Kesehatan. (2013). Riset Kesehatan Dasar (RISKESDAS) 2013. Laporan Nasional 2013, 1–384. <https://doi.org/1> Desember 2013

Becker, N. B., Isabel, R., Martins, S., Jesus, S. D. N., Chiodelli, R., & Stephen, M. (2018). Sleep health assessment : A scale validation, 259(August 2017), 51–55. <https://doi.org/10.1016/j.psychres.2017.10.014> diakses 14 September 2017 dari [http://www.psy-journal.com/article/S0165-1781\(17\)31545-7/fulltext](http://www.psy-journal.com/article/S0165-1781(17)31545-7/fulltext)

Black, J.M., & Hawks, J.H. (2009). *Medical surgical nursing: Clinical management for positive outcomes*. 8th Edition. Philadelphia: W.B. Sounders Company

Breivik, H., Borchgrevink, P. C., Allen, S. M., Rosseland, L. A., Romundstad, L., Breivik Hals, E. K., ... Stubhaug, A. (2008). Assessment of pain. *British Journal of Anaesthesia*, 101(1), 17–24. <https://doi.org/10.1093/bja/aen103> diakses 17 Januari 2017 dari [http://bjanaesthesia.org/article/S0007-0912\(17\)34263-0/pdf](http://bjanaesthesia.org/article/S0007-0912(17)34263-0/pdf)

Bulechek, G.M., Butcher, H & Dochterman, J M. 2013. *Nursing Intervention Classification (NIC)*. 6th edition. United States of America. Elsevier.

Burckhardt CS. The use of the McGill Pain Questionnaire in assessing arthritis pain. Pain 1984;19:305–14.

- Buysse, D., Reynold, C., Monk, T., Berman, S., Kupfer, D. 1989. The Pittsburgh Sleep Quality Index : A New Instrument for Psychiatric Practice and Research. *Psychiatry Res*; 28(2):193-213.
- Cappuccio, F.P., D'Elia, L., Strazzullo, P., and Miller, M.A. (2010). A systematic review and meta-analysis. Quantity and quality of sleep and incidence of type 2 diabetes. *Diabetes Care* (33) : 414–420, 2010 diakses 14 September 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/19910503>
- Caravaca, F., Gonzales, B., Bayo, M. A., & Luna, E. (2016). Musculoskeletal pain in patients with chronic kidney disease. *Nefrologia*, 36(4), 433–440. <https://doi.org/http://dx.doi.org/10.1016/j.nefro.2016.03.02> 4 diakses 14 September 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/27267921>
- Cengić, B., Resić, H., Spasovski, G., Avdić, E., & Alajbegović, A. (2012). Quality of sleep in patients undergoing hemodialysis. *International Urology and Nephrology*, 44(2), 557–67. <https://doi.org/10.1007/s11255-010-9881-x> diakses 14 September 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/21152979>
- Chang, Y., Cheng, S. Y., Lin, M., Gau, F. Y., & Chao, Y. F. C. (2010). The effectiveness of intradialytic leg ergometry exercise for improving sedentary life style and fatigue among patients with chronic kidney disease: A randomized clinical trial. *International Journal of Nursing Studies*, 47(11), 1383–1388. <https://doi.org/10.1016/j.ijnurstu.2010.05.002> diakses 14 September 2017 dari <https://doi.org/10.1016/j.ijnurstu.2010.05.002>
- Charalambous, A., Giannakopoulou, M., Bozas, E., Marcou, Y., Kitsios, P., & Paikousis, L. (2016). Guided imagery and progressive muscle relaxation as a cluster of symptoms management intervention in patients receiving

chemotherapy: A randomized control trial. *PLoS ONE*, 11(6), 1–18. <https://doi.org/10.1371/journal.pone.0156911>
diakses 14 September 2017
dari <https://www.ncbi.nlm.nih.gov/pubmed/27341675>

Conrad, A., & Roth, W. T. (2007). Muscle relaxation for anxiety disorder: It works but how?. *The Journal of Anxiety Disorder*, 22, 243-264 diakses 14 Juni 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/16949248>

Dahlan, Sopiyudin,, 2011. Statistik Untuk Kedokteran dan Kesehatan. Edisi 5.Jakarta: Salemba Medika

Davis, M. (2007). *The relaxation & stress reduction workbook bahasa Indonesia*. Penerjemah: Achiryan S Hamid dan Budi Anna Keliat. Jakarta: EGC

Davison, S. N., & Jhangri, G. S. (2010). Impact of Pain and Symptom Burden on the Health-Related Quality of Life of Hemodialysis Patients. *Journal of Pain and Symptom Management*, 39(3), 477–485. <https://doi.org/10.1016/j.jpainsymman.2009.08.008> diakses 17 Juli 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/20303025>

Dayapoğlu, N., & Tan, M. (2012). Evaluation of the effect of progressive relaxation exercises on fatigue and sleep quality in patients with multiple sclerosis. *Journal of Alternative and Complementary Medicine (New York, N.Y.)*, 18(10), 983–7. <https://doi.org/10.1089/acm.2011.0390> diakses 17 Juli 2017
dari <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3469207/>

Dewi SE, Petpitchetian W, Songwathana P. 2007. Comparison of pain experiences between Javanese and Batak patients undergoing major surgery in Medan, Indonesia. *SongklaNagarind Medical Journal* ; 25(4): 247-58.

Dharma, Kusuma Kelana. (2011). *Metodologi Penelitian Keperawatan : Panduan Melaksanakan dan Menerapkan Hasil Penelitian*. Jakarta: Trans InfoMedia

Duma (2012). *Pengaruh progressive muscle relaxation dan logoterapi terhadap ansietas dan depresi, kemampuan relaksasi dan kemampuan memaknai hidup klien kanker di RS Kanker Dharmais*. Universitas Indonesia. Jakartadiakses 17 Juli 2017 dari <http://lontar.ui.ac.id/file?file=digital/20305893-T30948%20-%20Pengaruh%20progressive.pdf>

Einollahi, B., Motalebi, M., Rostami, Z., Nemati, E., & Salesi, M. (2015). Sleep Quality Among Iranian Hemodialysis: A Multicenter Study. *Nephro-Urology Monthly*, 7(1), 1–7. <https://doi.org/10.5812/numonthly.23849> diakses 17 Juli 2017 dari <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4330687/>

Escalante A, Lichtenstein MJ, White K, Rios N, Hazuda HP. A method for scoring the pain map of the McGill pain questionnaire for use in epidemiologic studies. *Aging Clin Exp Res* 1995;7:358–66.

Elizabeth, Lindley, Aspinal, Claire & Garthwaite.(2011). Management Of Fluid Status In Haemodialysis Patients: The Roles Of Technologi And Dietary Advice. Departemen Of Renal Medicine, *Leeds Teaching Hospital NHS Trust United Kingdom*

Fonsêca, N. T., Santos, I. R., Fernandes, V., Thomaz Fernandes, V. A., Delgado Lopes, V. C., & Oliveira, L. V. F. (2014). Excessive daytime sleepiness in patients with chronic kidney disease undergone hemodialysis. *Fisioterapia Em Movimento*, 27(4), 653–660 8p. <https://doi.org/10.1590/0103-5150.027.004.AO17> diakses 17 Juli 2017 dari <http://www.scielo.br/pdf/fm/v27n4/0103-5150-fm-27-04-0653.pdf>

Ghonemy, T.A., Halla M. Allam, Amir M. Elokely, Yosef A. Kadry, Hesham M. Omar. 2016. Chronic pain in hemodialysis patients: Role of bone mineral metabolism. *Alexandria Journal of Medicine* (2016) 52, 337-342. Alexandria University Faculty of Medicine.

Greenberg, J. S. (2002). *Comprehensive stress management*. New York: Mc Graw-Hill Inc

Goldfarb, A. H. (n.d.). Exercise and Endogenous Opiates, 21–37. <https://doi.org/10.1007/978-1-62703-314-5>

Goumenos, D. S., Papachristou, E., & Papasotiriou, M. (2016). Renal replacement therapy in patients with heart and kidney failure. *Section of Medical Science*. Volume 37, Issue 2-3 <https://doi.org/10.1515/prilozi-2016-0015> diakses 18 Oktober 2017 dari <https://www.degruyter.com/downloadpdf/j/prilozi.2016.37.issue-2-3/prilozi-2016-0015/prilozi-2016-0015.pdf>

Harmono. (2010). *Pengaruh relaksasi otot progresif terhadap penurunan tekanan darah pada pasien hipertensi primer di Kota Malang*. Universitas Indonesia. Jakarta diakses 11 Juni 2017 dari lib.ui.ac.id/file?file=digital/20285357-T%20Rudi%20Harmono.pdf

Hastono, S.P. (2007). *Basic data analysis for health research training*. Depok: FKM UI

Hawker, G. A., Mian, S., Kendzerska, T., & French, M.(2011). Measures of adult pain: Visual analog scale for pain (VAS pain), numeric rating scale for pain (NRS pain), McGill pain questionnaire (MPQ), short-form McGill pain questionnaire (SF-MPQ), Chronic Pain Grade Scale (CPGS), short-form 36 bodily pain scale (SF-36 BPS), and measure of intermittent and constant osteoarthritis pain (ICOAP). *Arthritis Care and Research*, 63, 240–252

Hicks, C. L., Baeyer, C. L. Von, Spafford, P. A., Korlaar, I. Van, & Goodenough, B. (2001). The Faces Pain Scale ±

Revised : toward a common metric in pediatric pain measurement q, 93, 173–183. diakses 20 Desember 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/11427329>

Hill, N. R., Fatoba, S. T., Oke, J. L., Hirst, J. A., Callaghan, A. O., Lasserson, D. S., & Hobbs, F. D. R. (2016). Global Prevalence of Chronic Kidney Disease – A Systematic Review and Meta-Analysis, 1–18. <https://doi.org/10.5061/dryad.3s7rd>.Funding di akses 06 Juni 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/27383068>

Hjermstad, M. J., Fayers, P. M., Haugen, D. F., Caraceni, A., Hanks, G. W., Med, D., ... Kaasa, S. (2011). Studies Comparing Numerical Rating Scales , Verbal Rating Scales , and Visual Analogue Scales for Assessment of Pain Intensity in Adults : A Systematic Literature Review. *Journal of Pain and Symptom Management*, 41(6), 1073–1093. <https://doi.org/10.1016/j.jpainsymman.2010.08.016> diakses 07 Juli 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/21621130>

Hou, Y., Hu, P., & Liang, Y. (2014). Effects of Cognitive Behavioral Therapy on Insomnia of Maintenance Hemodialysis Patients. <https://doi.org/10.1007/s12013-014-9828-4> diakses 07 Juli 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/24577747>

Huang, T. H., Lin, S. C., Chang, F. L., Hsieh, S. S., Liu, S. H., & Yang, R. S. (2003). Effects of different exercise modes on mineralization, structure, and biomechanical properties of growing bone. *Journal of applied physiology*, 95(1), 300-307.

International Association for the Study of Pain. Visceral pain In : Core Curriculum for Professional Education in Pain. *IASP Press*. 2012; 34:1-2

Indonesian renal Registry. (2012). 5 th Report Of Indonesian Renal Registry 2012. *Program Indonesia Renal Registry*,

12–13. Retrieved from <http://www.pernefri-inasn.org/Laporan/5th Annual Report Of IRR 2012.pdf>

Indonesian Renal Registry. (2015). 7th Report Of Indonesian Renal Registry 2014. Program Indonesia Renal Registry. diakses 07 Juli 2017 dari <http://www.indonesianrenalregistry.org/data/INDONESIANRENALREGISTRY2014.pdf>

João, K.A.D.R., Becker, N.B., Jesus, S.N., Martins, R.I.S., 2017. Validation of the Portugal version of the Pittsburgh sleep quality index. *Psychiatry Res.* 247, 225–229. diakses 20 Desember 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/27923147>

Kallenbach, J.Z., Gutch, C.F., Martha, S.H., & Corca, A.L. (2005). *Review of Hemodialysis for nurses and dialysis personnel*. 7th edition. St. Louis: Elsevier Mosby

Kaplow, R. & Hardin, S.R., 2007. *Critical Care Nursing: Synergy For Optimal Outcomes*, Dallas Texas: Jones & Bartlett Learning.

Knutson, K.L., Phelan, J., Paskow, M.J., Roach, A., Whiton, K., Langer, G., Hillygus, D.S., Mokrzycki, M., Broughton, W.A., Chokroverty, S., Lichstein, K.L., Weaver, T.E., Hirshkowitz, M., 2017. The national sleep foundation's sleep health index. *Sleep Health*. 3, 234–240. <http://dx.doi.org/10.1016/j.slehd.2017.05.011>. di akses 13 september 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/28709508>

Koncicki, H. M., Unruh, M., & Schell, J. O. (2017). Pain Management in CKD: A Guide for Nephrology Providers. *American Journal of Kidney Diseases*, 69(3), 451–460. <https://doi.org/10.1053/j.ajkd.2016.08.039> di akses 18 Oktober 2017 dari <https://www.sciencedirect.com/science/article/pii/S0272638616305595>

- Lee, E. J., Bhattacharya, J., Sohn, C., Verres, R., (2012). Monochord sounds and progressive muscle relaxation reduce anxiety and improve relaxation during chemotherapy: A pilot EEG study complementary theraphies in medicine. *Elsevier Health Journal*, 20, 409-416 diakses 29 Mei 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/23131371>
- LeMone, P., & Burke, K. 2008. Medical surgical nursing: critical thinking in client care (4 th ed). New Jersey: Pearson Prenntice Hall
- Lindner, A. V., Novak, M., Bohra, M., & Mucsi, I. (2015). Insomnia in Patients With Chronic Kidney Disease. *Seminars in Nephrology*, 35(4), 359–372. <https://doi.org/10.1016/j.semephrol.2015.06.007> diakses 28 Juli 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/26355254>
- Tobing LumbanDuma. 2012. *Pengaruh progressive muscle relaxation dan logoterapi terhadap perubahan ansietas, depresi, kemampuan relaksasi dan kemampuan memaknai hidup klien kanker di RS Dharmais Jakarta*. Tesis. Universitas Indonesia
- Marchk, I. T. (2011). *Master your sleep, proven method simplified*. USA: Bascom Hils Publish Group.
- Mashudi. (2011). *Pengaruh progressive muscle relaxation terhadap kadar glukosa darah pasien diabetes mellitus tipe II Di RSUD Raden Matter Jambi*. Universitas Indonesia. Jakarta
- Maung, S. C., El Sara, A., Chapman, C., Cohen, D., & Cukor, D.(2017). Sleep disorders and chronic kidney disease. *World Journal Of Nephrology*, 5(3), 224
- Melzack R. The short-form McGill Pain Questionnaire. *Pain* 1987;30: 191–7.

Morgan GE, Mikhail MS, Murray MJ. *Pain Management*. In : Morgan GE, editor. Clinical Anesthesiology, 4th ed. Lange Medical Books/McGraw-Hill ; 2006. p. 359-412.

Neuendorf, R., Wahbeh, H., Chamine, I., Yu, J., Hutchison, K., & Oken, B. S. (2015). The Effects of Mind-Body Interventions on Sleep Quality: A Systematic Review. *Evidence-Based Complementary and Alternative Medicine*, 2015. <https://doi.org/10.1155/2015/902708> diakses 27 Mei 2017 dari <https://www.hindawi.com/journals/ecam/2015/902708/>

Notoatmodjo, S. (2012). *Metodologi Penelitian Kesehatan*. Jakarta: Rineka Cipta.

Nursalam. (2013). *Konsep Penerapan Metode Penelitian Ilmu Keperawatan*. Jakarta: Salemba Medika

Nursalam. (2016). *Metodologi Penelitian Ilmu Keperawatan*. Jakarta: Salemba Medika

Osman, N. a., Hassanein, S. M., Leil, M. M., & NasrAllah, M. M. (2015). Complementary and Alternative Medicine Use Among Patients With Chronic Kidney Disease and Kidney Transplant Recipients. *Journal of Renal Nutrition*, 25(6), 466–471. <https://doi.org/10.1053/j.jrn.2015.04.009> diakses 25 Juni 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/26093884>

Otsuki, T. et al. (2016) ‘Efficacy and Safety of Pregabalin for the Treatment of Neuropathic Pain in Patients Undergoing Hemodialysis’, *Clinical Drug Investigation*. Springer International Publishing, 23, p. 23. diakses 25 Agustus 2017 dari doi: <http://dx.doi.org/10.1007/s40261-016-0464-1>.

Parvan, K., Lakdizaji, S., Roshangar, F., & Mostofi, M. (2013). Quality of sleep and its relationship to quality of life in hemodialysis patients. *Journal of Caring Sciences*, 2(4), 295–304. <https://doi.org/10.5681/jcs.2013.035> diakses 18

Mei 2017 dari <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4134147>

Pearce, E. C. (2008). *Anatomi dan Fisiologi Untuk Tenaga Medis*. Jakarta: PT. Gramedia Pustaka Utama

Pham, P. C., Khaing, K., Sievers, T. M., Pham, P. M., Miller, J. M., Pham, S. V, ... Pham, P. T. (2017). 2017 update on pain management in patients with chronic kidney disease. *Clin Kidney J*, 1–10. <https://doi.org/10.1093/ckj/sfx080> diakses 18 Oktober 2017 dari <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5622905/>

Polit DF & Back, CT. 2012. Nursing Research. *Generating and Assessing Evidence for Nursing Practice*. 9th ed. Philadelphia: JB. Lippincott.

Potter & Perry. (2010). *Fundamental keperawatan*. Edisi 7 Buku 2. Jakarta. Salemba Medika

Prastika, D. 2018. Pain experience, pain management strategies and satisfaction of hospitalized trauma patients in Indonesia. *Journal of Nursing Education and Practice*, 8(1), 79–86. <https://doi.org/10.5430/jnep.v8n1p79>

Prawitasari, J. E. (2011). Psikologi Klinis: Pengantar Terapan Mikro & Makro. Jakarta: Erlangga

Price Sylvia A,& Wilson Lorraine M. 2014. Patofisiologi: Konsep KlinisProses-Proses Penyakit. Jakarta: EGC

Ramdhani, N., & Putra, A.A. (2008). Pengembangan multi media relaksasi. *Laporan Penelitian*. Yogyakarta: Fakultas Psikologi UGMdiakses 17 Oktober 2017 dari <http://neila.staff.ugm.ac.id/wordpress/wpcontent/uploads/2008/05/relaksasi-otot.pdf>

Rawal N, Fischer HBJ, Ivani G, Andreas JD, Mogensen T, Narchi P. (2008). Post operative pain management – good

clinical practice. *European Society of Regional Anaesthesia*. Sweden.

Roehrs T, Roth T.(2008). Caffeine: sleep and daytime sleepiness. *Sleep Med Rev*. doi: 10.1016/j.smrv.2007.07.004 diakses tanggal 17 Agustus 2018 dari <https://www.ncbi.nlm.nih.gov/pubmed/17950009>

Rudy Gunawan. (2011). Perbandingan Efek Paracetamol 1 gr/6 jam Intravena dan Ketorolak 30 mg/6 jam Intravena untuk Penanganan Nyeri Paska PembedahanSeksio Sesaria dengan Anestesi Regional Blok Subarahnoid. Thesis: Universitas Sumatera Utara

Ida Rosdiana. 2011. *Analisis faktor yang berhubungan dengan kejadian insomnia pada pasien gagal ginjal kronik yang menjalani hemodialisis di rumah sakit umum daerah kota Tasikmalaya dan Garut*. Tesis. Universitas Indonesia

Saeedi, M., Ashktorab, T., Saatchi, K., Zayeri, F., Amir, S., & Akbari, A. (2012). The Effect of Progressive Muscle Relaxation on Sleep Quality of Patients Undergoing Hemodialysis, 5(1), 23–28.

Saeedi, M., Shamsikhani, S., Farahani, P. V., & Haghverdi, F. (2014). Sleep hygiene training program for patients on hemodialysis. *Iranian Journal of Kidney Diseases*, 8(1), 65–69. diakses 20 Juli 2017 dari <http://www.ijkd.org/index.php/ijkd/article/viewArticle/1135>

Santoro, D., Satta, E., Messina, S., Costantino, G., Savica, V., & Bellinghieri, G. (2012). Pain in end-stage renal disease: a frequent and neglected clinical problem. *Clinical Nephrology*, 79. <https://doi.org/10.5414/CNX77S104> diakses 27 Juli 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/23249527>

Scherer, J. S., Combs, S. A. and Brennan, F. (2017) ‘Sleep Disorders, Restless Legs Syndrome, and Uremic Pruritus: Diagnosis and Treatment of Common Symptoms in

Dialysis Patients', *American Journal of Kidney Diseases*. Elsevier Inc, 69(1), pp. 117–128. diakses 27 September 2018 dari doi: 10.1053/j.ajkd.2016.07.031.

Scott, D., & Davidson, J. A. (2011). Managing Chronic Kidney Disease in Type 2 Diabetes in Family Practice. *Journal Of The National Medical Association*, 952-959.

Shariati, A., Jahani, S., Hooshmand, M., & Khalili, N. (2012). The effect of acupressure on sleep quality in hemodialysis patients. *Complementary Therapies in Medicine*, 20(6), 417–423. <https://doi.org/10.1016/j.ctim.2012.08.001> diakses 18 Oktober 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/23131372>

Smeltzer, S. C., Bare, B., Hinkle, J. L., & Cheever, K. H. 2010. *Brunner & Suddarth's Textbook of Medical Surgical Nursing*. China: Lippincott Williams & Wilkins

Snyder, M & Lindquist, R. (2006). *Complementary/alternative terapies in nursing(fifth edition)*. New York: Springer Publishing Company

Sudoyo, A. (2010). *Buku Ajar Ilmu Penyakit Dalam*. Jakarta: Balai Penerbit FKUI.

Wiwit Sugiarti. (2017). *Efektifitas intradyalitic exercise terhadap penurunan nyeri dan insomnia pasien hemodialisa rutin di unit hemodialisa RSUD Dr. Tjitrowardojo Purworejo*. Tesis. Universitas Muhammadiyah Yogyakarta

Sugiyono. (2012). *Statistika Untuk Penelitian*. Alfabeta. Bandung.

Sukandar, E. (2011). *Nefrologi Klinik*, Edisi 3. *Pusat Informasi Ilmiah Bagian Ilmu Penyakit Dalam*. Bandung. FK UNPAD RS Hasan Sadikin

Sumida, K., Molnar, M. Z., Potukuchi, P. K., & Thomas, F. (2017). Constipation and Incident CKD, 1248–1258.

<https://doi.org/10.1681/ASN.2016060656> diakses 17 Desember 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/28122944>

Sun, J., Kang, J., Wang, P., & Zeng, H. (2013). Self-relaxation training can improve sleep quality and cognitive functions in the older: A one-year randomised controlled trial. *Journal of Clinical Nursing*, 22(9–10), 1270–1280. <https://doi.org/10.1111/jocn.12096> diakses 18 Desember 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/23574290>

Suwanto, Agus Wiwit.(2018). Efektifitas Relaksasi Benson Terhadap Penurunan Stres dan Peningkatan Kualitas Tidur pada Pasien Hemodialisa. Tesis. Universitas Muhammadiyah Yogyakarta

Tamsuri, A. (2007). Konsep Dan Penatalaksanaan Nyeri. Jakarta: EGC

Tanase-Nakao, K., Naruse, M., Nanba, K., Tsuiki, M., Tagami, T., Usui, T., Okuno, H., Shimatsu, A., Hashimoto, S., Katabami, T., Ogo, A., Okumura, A., Umakoshi, H., Suzuki, T.(2014). Chronic Kidney Disease Score for Predicting Postoperative Masked Renal Insufficiency in Patients with Primary Aldosteronism. *Clinical Endocrinol*, 81(5):665-70. doi: 10.1111/cen.12497. diakses 16 Juni 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/24821606>

Teles, FD. (2014). Depression in Hemodialysis patients: the role of dialysis shift. *CLINICS*;69(3), 198-202

Thomas, N. (2010). *Renal Nursing*. 2nd Edition. London United Kingdom: Elsivier Science

Tohme, F. et al. (2017) ‘Predictors and outcomes of non-adherence in patients receiving maintenance hemodialysis’, *International Urology and Nephrology*. Springer Netherlands. diakses 17 September 2017 dari doi: 10.1007/s11255-017-1600-4.

- Tomb, A.D (2012). Buku saku psikiatri. Edisi 6. Jakarta: EGC
- Turkmen, K., Yazici, R., Solak, Y., Guney, I., Altintepe, L., Yeksan, M., & Tonbul, H. Z. (2012). Health-related quality of life, sleep quality, and depression in peritoneal dialysis and hemodialysis patients. *Hemodialysis International*, 16(2), 198–206. <https://doi.org/10.1111/j.1542-4758.2011.00648.x> diakses 17 Juni 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/22136456>
- Tzu. (2010). Evaluating integrated healthcare delivery models: A holistic approach. *Focus on Alternative and Complementary Therapy*, 7(4), 330-333. doi: 10.1211/fact.2002.00434
- United States Renal Data System. (2015). 2015 USRDS Annual Data Report. *United States Renal Data System*, 2(1), 227–238. <https://doi.org/10.1053/j.ajkd.2015.04.017>
- Vanholder, R., Fouque, D., Glorieux, G., Heine, G. H., Kanbay, M., Mallamaci, F., ... Lyon, C. B. (2016). Clinical management of the uraemic syndrome in chronic kidney disease. *THE LANCET Diabetes & Endocrinology*, 1–14. [https://doi.org/10.1016/S2213-8587\(16\)00033-4](https://doi.org/10.1016/S2213-8587(16)00033-4) diakses 18 Desember 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/26948372>
- Webster LR. (2015). Opioid-induced constipation. *Pain Med*. 2015;16(suppl 1):S16-S21.10 diakses 18 Mei 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/26461071>
- Weisbord S, Mor M, Green J, Sevick M, et. al. (2013). Comparison of symptom management strategies for pain, erectile dysfunction and depression in patients receiving chronic hemodialysis: A cluster randomized effectiveness trial. *Clinical Journal of the American Society of Nephrology*, vol. 8 (1) pp.90-99. Doi: 10.2215/CJN.044450512 di akses 20 Agustus 2018 dari <https://www.ncbi.nlm.nih.gov/pubmed/23024159>

Yom YH & Lee KE. (2008). A comparison of the knowledge of, experience with and attitudes towards complementary and alternative medicine between nurses and patients in Korea. *Journal of Clinical Nursing* 17, 2565– 2572. diakses 19 Mei 2017 dari <http://onlinelibrary.wiley.com/doi/10.1111/j.1365-2702.2007.02065.x/abstract>

Zyga, S., Alikari, V., Sachlas, A., Stathoulis, J., Aroni, A., Theofilou, P., & Panoutsopoulos, G. (2015). Management of Pain and Quality of Life in Patients with Chronic Kidney Disease Undergoing Hemodialysis. *Pain Management Nursing*, 16(5), 712–720. <https://doi.org/10.1016/j.pmn.2015.03.004> diakses 20 September 2017 dari <https://www.ncbi.nlm.nih.gov/pubmed/26162558>