

DAFTAR PUSTAKA

- American Diabetes Association. (2017). Standards of medical care in diabetes-2015. *The Journal Of Clinical And Applied Research And Education*, Vol.40, Sup.1
- Azab B, Daoud J, Naeem FB, Nasr R, Ross J, Ghimire P, Siddiqui A, Azzi N, Rihana N, Abdallah M, Azzi N, Patel P, Kleiner M, El-Sayegh S. (2012). *Neutrophil-to-Lymphocyte Ratio as a predictor of worsening renal function in diabetic patients (3-year follow-up study)*, In: *Renal Failure*, Vol. 34, hal.571-5767
- Anonym. Klinik Diabetes Nusantara (2007). Kontrol HbA1c.
- Buraerah, Hakim. Analisis Faktor Risiko Diabetes Melitus tipe 2 di Puskesmas Tanrutedong, Sidenreg Rappan,. *Jurnal Ilmiah Nasional*;2010 [cited 2010 feb 17].
- Baradero, Mary., Dayrit, Wilfrid, Mary., & Siswadi, Yakobus. (2009). *Seri Asuhan Keperawatan Klien Gangguan Endokrin*. Jakarta: EGC
- Chawla A, Chawla R, Jaggi S. Microvascular and macrovascular complications in diabetes mellitus: Distinct or continuum?. *Indian J Endocr Metab* 2016;20:546-53.
- Chittawar S, Dutta D, Qureshi Z, Surana V, Khandare S, Dubey TN. Neutrophil-lymphocyte ratio is a novel reliable predictor of nephropathy, retinopathy, and coronary artery disease in Indians with type-2 diabetes. *Indian J Endocr Metab* 2017;21:864-70.
- Darryl, R. Meeking: *Diabetes & Endocrinology*. 2011. Manson Publishing
- Depkes RI. (2008). *Riset Kesehatan Dasar Tahun 2008*. Badan Penelitian dan Pengembangan Kesehatan Departemen Kesehatan RI
- Dinas Kesehatan Kota Yogyakarta (2015). *Survailans Terpadu Penyakit (STP) Puskesmas dan Sistem Informasi Rumah Sakit (SIRS) di Provinsi DIY pada tahun 2015*.
- Fatih Sefil, Kemal Turker Ulutas. Recep Dokuyucu , Ahmet Taner Sumbul, Erhan Yengil, Abdullah Erman Yagiz, Erkan Yula, Ihsan Ustun and Cumali Gokce. *Investigation of neutrophil lymphocyte ratio and blood glucose regulation in patients with type 2 diabetes mellitus*. 2014, Vol. 42(2) 581–588. *Journal of International Medical Research*

- Fatimah, RN. Diabetes Melitus tipe 2. *Review article*. 2015. Vol.4, No.5. J Majority
- Forget Patrice, Khalifa Celine, Defour Jean-Philippe, Latinne Dominique, Van Pel Marie-Cecile, dan De Kock Marc. (2017). *What is the normal value of the neutrophil-to-lymphocyte ratio?*, Vol.10, No.12, hal.1-4
- Gokularishnan K, Deepa R, Mohan V. (2008). *Association of high sensitivity C-Reactive Protein [hsCRP] and Tumour Necrosis Factor- α [TNF- α] with carotid Intimal Medial Thickness in subjects with different grades of glucose intolerance—The Chennai Urban Rural Epidemiology Study (CURES-31)*, Vol.41, hal.480-485
- Gundogan K, Bayram F, Capak M, et al. Prevalence of metabolic syndrome in the Mediterranean region of Turkey: evaluation of hypertension, diabetes mellitus, obesity, and dyslipidemia. *Metab Syndr Relat Disord* 2009; 7: 427–434.
- H a n L i , Xiangxue Lu, Ruifang Xiong, and Shixiang Wang. High Neutrophil-to Lymphocyte Ratio Predicts Cardiovascular Mortality in Chronic Hemodialysis Patients. Volume 2017, Article ID 9327136, 8 pages. Hindawi Publishing Corporation Mediators of Inflammation.
- Hussain M, Babar MZM, Akhtar L, Hussain MS. Neutrophil lymphocyte ratio (NLR): A well assessment tool of glycemic control in Type2 diabetic patients. *Pak J Med Sci*. 2017;33(6):1366-137.
- James, NMD. (2011). *Managing Diabetes with blood glucose control*.
- Janis, Indra. (2008). *Konversi BTA pada Penderita TB Paru Kategori 1 yang mendapat terapi Intensif dengan Diabetes Melitus Terkontrol dan Diabetes Melitus Tidak Terkontrol*.
- Manaf, A. (2014). *Insulin:mekanisme sekresi dan aspek metabolisme In Ilmu Penyakit Dalam 6thn Edition (Hal. 2350-2357). Jakarta: Interna Publishing*
- M. Erturk, H. A. Cakmak, O. Surgit et al., “The predictive value of elevated neutrophil to lymphocyte ratio for longterm cardiovascular mortality in peripheral arterial occlusive disease,” *Journal of Cardiology*, vol. 64, no. 5, pp. 371–376, 2014.
- Nayak, B. S., Sobrian, A., Latiff, K., Pope, D., Rampersad, A., Lourenço, K., & Samuel, N. (2014). The association of age, gender, ethnicity, family history, obesity and hypertension with type 2 diabetes mellitus in Trinidad. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 8(2), 91–95.

- Ohshita K, Yamane K, Hanafusa M, et al. Elevated white blood cell count in subjects with impaired glucose tolerance. *Diabetes Care* 2004; 27: 491–496.
- Oyumlu M, Yildis A, Yuksel M, Polat N, Bilik MZ, Akyuz A, Aydin A, Aydin M, Aceet H, Soydinc S. Platelet to lymphocyte ratio is a predictor of in hospital mortality patients with acute coronary syndrome. 2015 Apr; 15 (4): 277-83.
- Papa A, Emdin M, Passino C, et al. Predictive value of elevated neutrophil-lymphocyte ratio on cardiac mortality in patients with stable coronary artery disease. *Clin Chim Acta* 2008; 395: 27–31.
- Putu Ugi Sugandha, AA Wiradewi Lestari. (2014). *Gambaran Pengendalian Kadar Gula Darah dan HbA1c pada pasien diabetes mellitus tipe 2 yang dirawat di RSUP Sanglah periode Januari- Mei 2014*. Denpasar.
- PERKENI. (2015). *Konsensus Pengelolaan dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia*. Jakarta:PERKENI
- Powers AC. Diabetes mellitus. *In: Fauci AS, Braunwald E, Kasper DL, Hauser SL, Longo DL, Jameson JL. Harrison's Principles of Internal Medicine. 19th Edition. United States: The McGraw-Hill Companies; 2015, hal.2399-2422.*
- Riskesmas (2013). *Riset Kesehatan Dasar Tahun 2013*. Badan Penelitian dan Pengembangan Kesehatan Departemen Kesehatan RI
- Soegondo, S. (2009). *Sindrom Metabolik In Ilmu penyakit dalam (5th ed) (hal.1865- 1872)*. Jakarta: Interna Publishing
- Sterne, J. Treatment of diabetes mellitus with N,Ndimethylguanylguanidine (LA.6023, glucophage). *Therapie* 1959, Vol.14, hal.625-630
- Stratton IM, Adler AI, Neil HA, Matthews DR, Manley SE, Cull CA, Hadden D, Turner RC, Holman RR. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ*. 2000 Aug 12; 321(7258): 405-412
- Tsai JC, Sheu SH, Chiu HC, et al. Association of peripheral total and differential leukocyte counts with metabolic syndrome and risk of ischemic cardiovascular diseases in patients with type 2 diabetes mellitus. *Diabetes Metab Res Rev* 2007; 23: 111–118.
- Wandell PE dan Carlsson AC. (2014). Gender differences and time trends in incidence and prevalence of type 2 diabetes in Sweden—a model explaining the diabetes epidemic worldwide today? *Diabetes Res Clin Pract*. Vol.106, hal.90-92.
- World Health Organization (2016). *Global Report on Diabetes*. Geneva: diakses dari (http://www.who.int/about/licensing/copyright_form/index.html)