

LAMPIRAN 1. TABEL STATISTIK

Karakteristik Responden

Umur

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid ≤ 55 Tahun	26	26.3	26.3	26.3
> 55 Tahun	73	73.7	73.7	100.0
Total	99	100.0	100.0	

Lama Rawat

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid < 7 Hari	73	73.7	73.7	73.7
≥7 Hari	26	26.3	26.3	100.0
Total	99	100.0	100.0	

Kadar LDL

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Normal	52	52.5	52.5	52.5
Tinggi	47	47.5	47.5	100.0
Total	99	100.0	100.0	

Kadar Kolesterol Total

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Normal	65	65.7	65.7	65.7
Tinggi	34	34.3	34.3	100.0
Total	99	100.0	100.0	

Gula Darah

	Frequency	Percent	Valid Percent	Cumulative Percent
--	-----------	---------	---------------	--------------------

	Normal	85	85.9	85.9	85.9
Valid	Tinggi	14	14.1	14.1	100.0
	Total	99	100.0	100.0	

Tekanan Darah

		Frequency	Percent	Valid Percent	Cumulative Percent
	Normal	18	18.2	18.2	18.2
Valid	Tinggi	81	81.8	81.8	100.0
	Total	99	100.0	100.0	

Uji Bivariat

Kadar LDL * Lama Rawat

Crosstab

			Lama Rawat		Total
			< 7 Hari	≥7 Hari	
Kadar LDL	Normal	Count	35	17	52
		% of Total	35.4%	17.2%	52.5%
	Tinggi	Count	38	9	47
		% of Total	38.4%	9.1%	47.5%
Total	Count	73	26	99	
	% of Total	73.7%	26.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	2.338 ^a	1	.126		
Continuity Correction ^b	1.691	1	.193		
Likelihood Ratio	2.373	1	.123		
Fisher's Exact Test				.171	.096
Linear-by-Linear Association	2.315	1	.128		
N of Valid Cases	99				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 12.34.

b. Computed only for a 2x2 table

Kadar Kolesterol Total * Lama Rawat

Crosstab

			Lama Rawat		Total
			< 7 Hari	≥7 Hari	
Kadar Kolesterol Total	Normal	Count	54	11	65
		% of Total	54.5%	11.1%	65.7%
	Tinggi	Count	19	15	34
		% of Total	19.2%	15.2%	34.3%
Total	Count	73	26	99	
	% of Total	73.7%	26.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	8.525 ^a	1	.004		
Continuity Correction ^b	7.179	1	.007		
Likelihood Ratio	8.237	1	.004		
Fisher's Exact Test				.007	.004
Linear-by-Linear Association	8.439	1	.004		
N of Valid Cases	99				

a. 0 cells (0.0%) have expected count less than 5. The minimum expected count is 8.93.

b. Computed only for a 2x2 table

Hiperglikemia * Lama Rawat

Crosstab

			Lama Rawat		Total
			< 7 Hari	≥7 Hari	
Hiperglikemia	Normal	Count	60	25	85
		% of Total	60.6%	25.3%	85.9%
	Tinggi	Count	13	1	14
		% of Total	13.1%	1.0%	14.1%
Total	Count	73	26	99	
	% of Total	73.7%	26.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	3.078 ^a	1	.079		
Continuity Correction ^b	2.036	1	.154		
Likelihood Ratio	3.815	1	.051		
Fisher's Exact Test				.105	.069
Linear-by-Linear Association	3.047	1	.081		
N of Valid Cases	99				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 3.68.

b. Computed only for a 2x2 table

Tekanan Darah * Lama Rawat

Crosstab

			Lama Rawat		Total
			< 7 Hari	≥7 Hari	
Tekanan Darah	Normal	Count	13	5	18
		% of Total	13.1%	5.1%	18.2%
	Tinggi	Count	60	21	81
		% of Total	60.6%	21.2%	81.8%
Total	Count	73	26	99	
	% of Total	73.7%	26.3%	100.0%	

Chi-Square Tests

	Value	df	Asymp. Sig. (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.026 ^a	1	.872		
Continuity Correction ^b	.000	1	1.000		
Likelihood Ratio	.026	1	.872		
Fisher's Exact Test				1.000	.541
Linear-by-Linear Association	.026	1	.872		
N of Valid Cases	99				

a. 1 cells (25.0%) have expected count less than 5. The minimum expected count is 4.73.

b. Computed only for a 2x2 table