

## DAFTAR PUSTAKA

- Agarwal, R. (2016) 'Defining end-stage renal disease in clinical trials: a framework for adjudication: Table 1.', *Nephrology Dialysis Transplantation*, 31(6), pp. 864–867. doi: 10.1093/ndt/gfv289.
- Al-Rubeaan, K. et al. (2015) 'Diabetic retinopathy and its risk factors in a society with a type 2 diabetes epidemic: A Saudi National Diabetes Registry-based study', *Acta Ophthalmologica*, 93(2), pp. e140–e147. doi: 10.1111/aos.12532.
- Alrawahi, A. H. et al. (2012) 'Prevalence and risk factors of diabetic nephropathy in Omani type 2 diabetics in Al-Dakhiliyah region', *Oman Medical Journal*, 27(3), pp. 212–216. doi: 10.5001/omj.2012.48.
- American Diabetes Association (2018) 'STANDARDS OF MEDICAL CARE IN DIABETES — 2018', 41(January).
- Atapour, A. et al. (2015) 'Hemodialysis Versus Peritoneal Dialysis, Which is Cost-effective?', *Saudi Journal of Kidney Diseases and Transplantation*, 26(5), pp. 962–965.
- Bennett, K. and Aditya, B. S. (2015) 'An overview of diabetic nephropathy: Epidemiology, pathophysiology and treatment', *Journal of Diabetes Nursing*, 19(2), pp. 61–67.
- Clotet, S. et al. (2016) 'RAS and sex differences in diabetic nephropathy Sergi', *Am J Physiol Renal Physiol*, pp. F945–F957. doi: 10.1152/ajprenal.00292.2015.
- Defronzo, R. A. (2009) 'From the triumvirate to the ominous octet: A new paradigm for the treatment of type 2 diabetes mellitus', *Diabetes*, 58(4), pp. 773–795. doi: 10.2337/db09-9028.
- Fowler, M. J. (2014) 'Microvascular and Macrovascular Complications of Diabetes', 26(2), pp. 77–82. doi: <http://dx.doi.org/10.2337/diaclin.26.2.77>.
- François, K., Ronco, C. and Bargman, J. M. (2015) 'Peritoneal Dialysis for Chronic Congestive Heart Failure', *Blood Purification*, 40(1), pp. 45–52. doi: 10.1159/000430084.
- Galloway, D. A. et al. (2015) 'What Is Known and Unknown About Twice-Weekly Hemodialysis', *HHS Public Access*, 40(4), pp. 298–305. doi: 10.1016/j.coviro.2015.09.001.Human.
- Ghaderian, S. B. et al. (2015) 'Diabetes and end-stage renal disease; a review article on new concepts.', *Journal of renal injury prevention*, 4(2), pp. 28–33. doi: 10.12861/jrip.2015.07.
- Halle, M. P. et al. (2015) 'Epidemiological profile of patients with end stage renal disease in a referral hospital in Cameroon', *BMC Nephrology*, 16(1), p. 59. doi: 10.1186/s12882-015-0044-2.
- Huang, G.-M. et al. (2015) 'An interpretable rule-based diagnostic classification of diabetic nephropathy among type 2 diabetes patients', *BMC Bioinformatics*. BioMed Central Ltd, 16(Suppl 1), p. S5. doi: 10.1186/1471-2105-16-S1-S5.
- Intensive Care Society (2009) 'Standards and Recommendations for the Provision of Renal Replacement Therapy on Intensive Care Units in the United Kingdom', pp. 1–32.

- International Diabetes Federation (2017) *IDF Diabetes Atlas*. Available at: <http://www.diabetesatlas.org/resources/2017-atlas.html>.
- Joannidis, M. and Forni, L. G. (2011) 'Timing of Renal Replacement Therapy', *Critical Care*, 15(3), p. 223. doi: 10.1186/cc10109.
- Kemenkes RI (2013) 'Riset Kesehatan Dasar (RISKESDAS) 2013', *Laporan Nasional 2013*, pp. 1–384. doi: 1 Desember 2013.
- Lee, S. and Choi, M. E. (2016) 'Urinary biomarkers for early diabetic nephropathy: Beyond albuminuria', 30(7), pp. 1063–1075. doi: 10.1007/s00467-014-2888-2.Urinary.
- Lim, A. K. H. (2014) 'Diabetic nephropathy – Complications and treatment', *International Journal of Nephrology and Renovascular Disease*, 7, pp. 361–381. doi: 10.2147/IJNRD.S40172.
- Mota, E. et al. (2009) 'Risk factors for diabetic nephropathy: intrinsic or extrinsic renal?', *Rom J Intern Med*, 47(4), pp. 397–401. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21179923>.
- Muhammad, C. and Nazar, J. (2014) 'Diabetic nephropathy; principles of diagnosis and treatment of diabetic kidney disease', 3(1), pp. 15–20.
- National Kidney Foundation (2015) *K/DOQI Clinical Practice Guidelines for Chronic Kidney Disease: Evaluation, Classification and Stratification*, Am J Kidney Dis. doi: 10.1634/theoncologist.2011-S2-45.
- Nurjanah, A. (2012) 'Hubungan Antara Lama Hipertensi dengan Angka Kejadian Gagal Ginjal Terminal di RSUD dr. Moewardi Surakarta', pp. 1–15.
- Oudit, GY. et al. (2010) 'Human recombinant ACE2 reduces the progression of diabetic nephropathy', *Diabetes* 59: 529–538, 2010.
- PERKENI (2015) *Konsensus Pengendalian dan Pencegahan Diabetes Melitus Tipe 2 di Indonesia 2015*, Perkeni. doi: 10.1017/CBO9781107415324.004.
- Putra, I. W. A. et al. (2015) 'Empat Pilar Penatalaksanaan Pasien Diabetes Mellitus Tipe 2 Four Pillars of Management of Type 2 Diabetes Mellitus Patients', 4(Dm), pp. 8–12.
- Putri, R. I. (2014) 'FAKTOR DETERMINAN NEFROPATI DIABETIK PADA PENDERITA DIABETES MELLITUS DI RSUD DR. M. SOEWANDHIE SURABAYA', pp. 109–121.
- Rivandi, J. and Yonata, A. (2015) 'Hubungan Diabetes Melitus Dengan Kejadian Gagal Ginjal Kronik', majority, 4(9), pp. 27–34. Available at: <http://jukeunila.com/wp-content/uploads/2016/02/5.pdf>.
- Sakti, D. H. H. (2016) 'Hubungan Faktor Risiko pada Diabetes Melitus Terhadap Derajat Neuropati Diabetik pada Kelompok PERSADIA RSI Amal Sehat Sragen ', (Iddm), pp. 6–26.
- Shah, A. and Kanaya, A. M. (2014) 'Diabetes and associated complications in the South Asian population', *Curr Cardiol Rep*, 16(5), p. 476. doi: 10.1007/s11886-014-0476-5.
- Spruijt-Metz, D. et al. (2014) 'Behavioral contributions to the pathogenesis of type 2 diabetes', *Current Diabetes Reports*, 14(4), p. 475. doi: 10.1007/s11892-014-0475-3.
- Tillou, X. et al. (2014) 'Prostatectomy for Localized Prostate Cancer to Prepare for Renal Transplantation in End-Stage Renal Disease Patients', *Annals of*

- Transplantation*, 19(1), pp. 569–575. doi: 10.12659/AOT.890908.
- Tziomalos, K. and Athyros, V. G. (2015) ‘Diabetic nephropathy: New risk factors and improvements in diagnosis’, *Review of Diabetic Studies*, 12(1–2), pp. 110–118. doi: 10.1900/RDS.2015.12.110.
- Vujičić, B. et al. (2012) ‘Diabetic Nephropathy’, *Pathophysiology and Complications of Diabetes Mellitus*, 4, pp. 71–96. doi: <http://dx.doi.org/10.5772/50115>.
- World Health Organization (2016) ‘Global Report on Diabetes Executive Summary’, *Www.Who.Int/Diabetes/Global Report*, pp. 4–4. doi: 10/ISBN 978 92 4 156525 7.
- Wu, Y. et al. (2014) ‘Risk factors contributing to type 2 diabetes and recent advances in the treatment and prevention’, *International journal of medical sciences*, 11(11), pp. 1185–1200. doi: 10.7150/ijms.10001.
- Yamaleyeva, LM. et al. (2012) ‘Differential regulation of circulating and renal ACE2 and ACE in hypertensive mRen2 Lewis rats with early-onset diabetes’, *Am J Physiol Renal Physiol* 302: F1374–F1384
- Yang, C.-W. et al. (2011) ‘Prevalence of diabetic nephropathy in primary care type 2 diabetic patients with hypertension: data from the Korean Epidemiology Study on Hypertension III (KEY III study)’, *Nephrology Dialysis Transplantation*, 26(10), pp. 3249–3255. doi: 10.1093/ndt/gfr011.