

LAMPIRAN

KUESIONER PENELITIAN

HUBUNGAN INDEKS MASA TUBUH (IMT) PASIEN OSTEOARTRITIS LUTUT DITINJAU DARI PEMERIKSAAN RADIOLOGI TAHUN 2016

1. FORMULIR IDENTITAS RESPONDEN

Kode Responden :

Kelas :

Alamat :

Jenis kelamin :

Tanggal Lahir :

Tanggal pengukuran :

Umur :

2. DATA ANTROPOMETRI

Tinggi badan : cm

Berat badan : kg

$$\text{IMT} = \frac{\text{Massa Tubuh (kg)}}{\text{Tinggi Badan (m)} \times \text{Tinggi Badan (m)}}$$

Status Gizi	IMT (kg/m²)
Sangat Kurus	<14,9
Kurus	15,0-18,4
Normal	18,5-22,9
Massa Tubuh Berlebih	23,0-27,4
Obesitas	27,5-40,0
Sangat Obesitas	>40,0

*berdasarkan BMI Cuts-off in Singapore (2005)

**berlaku untuk dewasa berusia diatas 20 tahun

		Jenis_Kelamin	Usia	IMT
N	Valid	63	63	63
	Missing	0	0	0
Mean		1.6349	2.2857	4.2381
Median		2.0000	2.0000	4.0000
Mode		2.00	3.00	5.00
Std. Deviation		.48532	.72798	.79746
Minimum		1.00	1.00	2.00
Maximum		2.00	3.00	5.00

Jenis_Kelamin

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Laki-laki	23	36.5	36.5	36.5
	Perempuan	40	63.5	63.5	100.0
Total		63	100.0	100.0	

Usia

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	< 50 tahun	10	15.9	15.9	15.9
	50 - 60 tahun	25	39.7	39.7	55.6
	> 60 tahun	28	44.4	44.4	100.0
Total		63	100.0	100.0	

IMT

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid Kurus	1	1.6	1.6	1.6
Normal	11	17.5	17.5	19.0
Massa Tubuh Berlebih	23	36.5	36.5	55.6
Obesitas	28	44.4	44.4	100.0
Total	63	100.0	100.0	

Spesifikasi_Gmbaran_Radiologis_OA_Lutut

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Penyempitan celah sendi asimetris	27	38.0	38.0	38.0
	Peningkatan densitas sklerotik	19	26.8	26.8	64.8
	Osteofit	18	25.4	25.4	90.1
	Perubahan struktur tulang	7	9.9	9.9	100.0
	Total	71	100.0	100.0	

Lokasi_Gambaran_Radiologis_Pasien_OA_Lutut

		Frequency	Percent	Valid Percent	Cumulative Percent
Valid	Genu Dextra	17	23.9	35.4	35.4
	Genu Sinistra	15	21.1	31.3	66.7
	Genu Dextra Sinistra	16	22.5	33.3	100.0
	Total	48	67.6	100.0	
Missing	System	23	32.4		
Total		71	100.0		

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
Peneliti_A * Peneliti_B	5	100.0%	0	.0%	5	100.0%

Peneliti_A * Peneliti_B Crosstabulation

			Peneliti_B		Total
			OA	Tidak OA	OA
Peneliti_A	OA	Count	1	0	1
		Expected Count	.6	.4	1.0
		% within Peneliti_A	100.0%	.0%	100.0%
		% within Peneliti_B	33.3%	.0%	20.0%
		% of Total	20.0%	.0%	20.0%
	Tidak OA	Count	2	2	4
		Expected Count	2.4	1.6	4.0
		% within Peneliti_A	50.0%	50.0%	100.0%
		% within Peneliti_B	66.7%	100.0%	80.0%
		% of Total	40.0%	40.0%	80.0%
Total	Count	3	2	5	
	Expected Count	3.0	2.0	5.0	
	% within Peneliti_A	60.0%	40.0%	100.0%	
	% within Peneliti_B	100.0%	100.0%	100.0%	
	% of Total	60.0%	40.0%	100.0%	

Symmetric Measures

		Value	Asymp. Std. Error(a)	Approx. T(b)	Approx. Sig.
Measure of Agreement	Kappa	.286	.274	.913	.361
N of Valid Cases		5			

a Not assuming the null hypothesis.

b Using the asymptotic standard error assuming the null hypothesis.

Crosstab

			OA Lutut		Total
			Ya	Tidak	
IMT	Normal (<23)	Count	4	8	12
		% within IMT	33.3%	66.7%	100.0%
		% within OA Lutut	8.3%	53.3%	19.0%
	Berlebih (23-27,4)	Count	18	5	23
		% within IMT	78.3%	21.7%	100.0%
		% within OA Lutut	37.5%	33.3%	36.5%
	Obesitas (>27,4)	Count	26	2	28
		% within IMT	92.9%	7.1%	100.0%
		% within OA Lutut	54.2%	13.3%	44.4%
Total	Count	48	15	63	
	% within IMT	76.2%	23.8%	100.0%	
	% within OA Lutut	100.0%	100.0%	100.0%	

Chi-Square Tests

	Value	Df	Asymp. Sig. (2-sided)
Pearson Chi-Square	16.492(a)	2	.000
Likelihood Ratio	15.387	2	.000
Linear-by-Linear Association	14.526	1	.000
N of Valid Cases	63		

a. 1 cells (16.7%) have expected count less than 5. The minimum expected count is 2.86.

Directional Measures

			Value
Nominal by Interval	Eta	IMT Dependent	.484
		OA Lutut Dependent	.512